arXiv:1204.0758v1 [math.PR] 3 Apr 2012

One-sided FKPP travelling waves in the context of homogenous fragmentation processes

Robert Knobloch*

March 2, 2013

Abstract

In this paper we introduce the one-sided FKPP equation in the context of homogenous fragmentation processes. More specifically, the main result of the present paper is concerned with the existence and uniqueness of one-sided FKPP travelling waves in this setting. Moreover, we prove some analytical properties of such travelling waves. Our techniques make use of killed fragmentation processes and an associated product martingale as well as various properties of spectrally negative Lévy processes. Furthermore, as a motivation, we devote a section to fragmentations with a finite dislocation measure, where we obtain a stronger result than for general fragmentation processes.

2010 Mathematics Subject Classification: 60J25, 60G09.

Keywords and phrases: FKPP equation, travelling wave, homogenous fragmentation process, killed fragmentation, product martingale, spectrally negative Lévy process.

1 Introduction

This paper deals with the one-sided FKPP travelling wave equation in the setting of conservative as well as dissipative homogenous fragmentation processes. In particular, we are concerned with the existence of solutions to this equation and we derive certain analytical properties of the solutions when they exist. The FKPP travelling wave equation in the context of fragmentations has a similar probabilistic interpretation as the classical FKPP travelling wave equation whose probabilistic interpretation is related to branching Brownian motions, see Section 4. In this respect we also refer to the recent paper [BHK11], where the two-sided FKPP travelling wave equation for conservative homogenous fragmentations was studied. It turns out that solutions in the setting of fragmentation processes have similar properties as their classical counterparts. However, the techniques of proving existence and uniqueness results are very different between the two cases, not least because the equations differ significantly. Indeed, whereas the classical FKPP travelling wave equation is a differential equation of second order, the FKPP travelling wave equation in our setting is an integro-differential equation of first order. This difference results from the non-diffusive behaviour of fragmentation processes and the more complicated jump structure of fragmentations in comparison with branching Brownian motions.

---

*Institut für Mathematik, FB 12, Goethe-Universität Frankfurt am Main, 60054 Frankfurt am Main, Germany
e-mail: knobloch@math.uni-frankfurt.de

In the context of homogenous fragmentation processes we prove the existence and uniqueness of one-sided travelling waves within a certain regime of wave speeds. More precisely, the problem we are concerned with in this paper can be roughly described as follows. Consider the integro-differential equation

$$cf'(x) + \int_{\mathcal{P}} \left(\prod_n f(x + \ln(|\pi_n|)) - f(x) \right) \mu(\mathrm{d}\pi) = 0$$

for certain $c > 0$ and all $x \in \mathbb{R}_0^+$, where the product is taken over all $n \in \mathbb{N}$ with $|\pi_n| > 0$. Here the space $\mathcal{P}$ is the space of partitions $(\pi_n)_{n\in\mathbb{N}}$ of $\mathbb{N}$ and μ is the so-called dislocation measure on $\mathcal{P}$. This notation is introduced in more detail in the next section. We are interested in solutions $f : \mathbb{R} \to [0, 1]$ of the above equation that satisfy

$$f|_{\mathbb{R}_0^+} \in C^1(\mathbb{R}_0^+, [0, 1]) \qquad \text{and} \qquad f|_{(-\infty,0)} \equiv 1$$

as well as the boundary condition

$$\lim_{x\to\infty} f(x) = 0.$$

Roughly speaking, the main result of this paper states that there is some constant $c_0 > 0$ such that there exists a unique solution of the above boundary value problem for every $c > c_0$ and there does not exist such a solution for any $c \leq c_0$. Our approach is based on using fragmentation processes with killing at an exponential barrier. These processes have been studied in [KK11] and we briefly describe the corresponding concepts below.

The outline of this paper is as follows. In the next section we give a brief introduction to homogenous fragmentation processes as well as appropriately killed versions of such fragmentations. Subsequently, in Section 3 we describe the set-up of our situation and we state our main results. Afterwards, in the fourth section we provide some motivation for the problems considered in this paper by explaining some related results that are known in the literature for fragmentation processes and branching Brownian motions respectively. In Section 5 we show how the existence and uniqueness of one-sided FKPP travelling waves for fragmentation processes can be obtained if the dislocation measure is finite. This provides some motivation for the existence and uniqueness result for general fragmentation processes this paper is mainly concerned with. The subsequent three sections are devoted to the proofs of our main results.

Throughout the present paper we use the terminology $f : \mathbb{R} \to [0, 1]$ as an abbreviation for functions $f : \mathbb{R} \cup \{-\infty\} \to [0, 1]$ with $f(-\infty) = 1$, and similarly we interpret $u : \mathbb{R}_0^+ \times \mathbb{R} \to [0, 1]$. Let us point out that we denote by $\mathbb{R}_0^+$ the nonnegative and by $\mathbb{R}^+$ the positive real numbers. In addition, we adopt the conventions $\ln(0) := -\infty$ as well as $\inf(\emptyset) := \infty$. The notation C^n, $n \in \mathbb{N}_0$, refers to the set of n-times continuously differentiable functions. The integral with respect to Lebesgue measure of a real-valued function f on a set $[s, t] \subseteq \mathbb{R}$ is denoted by $\int_{[s,t]} f(u)\mathrm{d}u$ and $\int_s^t f(u)\mathrm{d}u$ denotes the Riemann integral. The operators $\wedge$ and $\vee$ refer to the minimum and maximum respectively. Furthermore, we consider a probability space $(\Omega, \mathscr{F}, \mathbb{P})$ on which all the random objects are defined.

2 Killed homogenous fragmentation processes

In this section we provide a brief introduction to partition-valued fragmentation processes and we present the main tools that we need in the subsequent sections. In addition, we introduce some specific killing scheme for these processes. The advantage of partition-valued fragmentation process compared to so-called mass fragmentations is their explicit *genealogical structure* of blocks. This structure is crucial for the killing mechanism that we introduce below.

Regarding the state space of partition-valued fragmentation processes let $\mathcal{P}$ be the space of partitions $\pi = (\pi_n)_{n\in\mathbb{N}}$ of $\mathbb{N}$, where the blocks of π are ordered by their least element such that $\inf(\pi_i) < \inf(\pi_j)$ if $i < j$. For every $\pi \in \mathcal{P}$ we denote by $(|\pi|_n^{\downarrow})_{n\in\mathbb{N}}$ the decreasing reordering of the asymptotic frequencies of the blocks of π, cf. (2). Throughout this paper we consider a homogenous $\mathcal{P}$-valued fragmentation process $\Pi := (\Pi(t))_{t\in\mathbb{R}_0^+}$, where $\Pi(t) = (\Pi_n(t))_{n\in\mathbb{N}}$. Homogenous $\mathcal{P}$-valued fragmentations are exchangeable Markov processes that were introduced in [Ber01], see also [Ber06]. Further, let $\mathscr{F} := (\mathscr{F}_t)_{t\in\mathbb{R}_0^+}$ be the filtration generated by Π. Bertoin showed in [Ber01] that the distribution of Π is determined by some constant $c \in \mathbb{R}_0^+$ (the *rate of erosion* which describes the drift of Π) and a σ-finite measure ν (the so-called *dislocation measure* that indirectly describes the jumps of Π) on the infinite simplex

$$\mathcal{S} := \left\{ \mathbf{s} := (s_n)_{n\in\mathbb{N}} : s_1 \geq s_2 \ldots \geq 0, \sum_{n\in\mathbb{N}} s_n \leq 1 \right\},$$

such that $\nu(\{(1, 0, \cdots)\}) = 0$ and

$$\int_{\mathcal{S}} (1 - s_1)\nu(d\mathbf{s}) < \infty. \tag{1}$$

The process Π is said to be *conservative* if $\nu(\sum_{n\in\mathbb{N}} s_n < 1) = 0$, i.e. if there is no loss of mass by sudden dislocations, and *dissipative* otherwise. In this paper we allow for both of these cases.

Throughout this paper we assume that $c = 0$ *as well as* $\nu(\mathbf{s} \in \mathcal{S} : s_2 = 0) = 0$.
This enables us to resort to the results of [KK11], where the same assumptions are made.

Consider the exchangeable partition measure μ on $\mathcal{P}$ given by

$$\mu(d\pi) = \int_{\mathcal{S}} \varrho_{\mathbf{s}}(d\pi)\nu(d\mathbf{s}),$$

where $\varrho_{\mathbf{s}}$ is the law of Kingman's paint-box based on $\mathbf{s} \in \mathcal{S}$. Similarly to ν the measure μ describes the jumps of Π, though more directly, and is also referred to as *dislocation measure*. In [Ber01] Bertoin showed that the homogenous fragmentation process Π is characterised by a Poisson point process. More precisely, there exists a $\mathcal{P} \times \mathbb{N}$-valued Poisson point process $(\pi(t), \kappa(t))_{t\in\mathbb{R}_0^+}$ with characteristic measure $\mu \otimes \sharp$, where $\sharp$ denotes the counting measure on $\mathbb{N}$, such that Π changes state precisely at the times $t \in \mathbb{R}_0^+$ for which an atom $(\pi(t), \kappa(t))$ occurs in $(\mathcal{P} \setminus (\mathbb{N}, \emptyset, \ldots)) \times \mathbb{N}$. At such a time $t \in \mathbb{R}_0^+$ the sequence $\Pi(t)$ is obtained from $\Pi(t-)$ by replacing its $\kappa(t)$-th term, $\Pi_{\kappa(t)}(t-) \subseteq \mathbb{N}$, with the restricted partition $\pi(t)|_{\Pi_{\kappa(t)}(t-)}$ and reordering the terms such that the resulting partition of $\mathbb{N}$ is an element of $\mathcal{P}$. We denote the random jump times of Π, i.e. the times at which the abovementioned Poisson point process has an atom in $(\mathcal{P} \setminus (\mathbb{N}, \emptyset, \ldots)) \times \mathbb{N}$, by $(t_i)_{i\in\mathcal{I}}$, where the index set $\mathcal{I} \subseteq \mathbb{R}_0^+$ is countably infinite.

Moreover, by exchangeability, the limits

$$|\Pi_n(t)| := \lim_{k\to\infty} \frac{\sharp(\Pi_n(t) \cap \{1, \ldots, k\})}{k}, \tag{2}$$

referred to as *asymptotic frequencies*, exist $\mathbb{P}$-a.s. simultaneously for all $t \in \mathbb{R}_0^+$ and $n \in \mathbb{N}$. Let us point out that the concept of asymptotic frequencies provides us with a notion of *size* for the blocks of a $\mathcal{P}$-valued fragmentation process. In [Ber01] Bertoin showed that the process $(-\ln(|\Pi_1(t)|))_{t\in\mathbb{R}_0^+}$ is a killed subordinator, a fact we shall make use of below.

For the time being, let $x \in \mathbb{R}$. In this paper we are concerned with a specific procedure of killing of Π, see Figure 1(a), that was introduced in [KK11]. More precisely, we kill Π upon hitting the space-time barrier

$$\left\{(y,t) \in \mathbb{R}_0^+ \times \mathbb{R}_0^+ : y < e^{-(x+ct)}\right\}.$$

That is to say, a block $\Pi_n(t)$ is killed at the moment of its creation $t \in \mathbb{R}_0^+$ if $|\Pi_n(t)| < e^{-(x+ct)}$. We denote the resulting killed fragmentation process, with cemetery state $(\emptyset, \ldots)$, by $\Pi^x := (\Pi^x(t))_{t\in\mathbb{R}_0^+}$. For any $t \in \mathbb{R}_0^+$ let $\mathcal{N}_t^x$ denote the set of indices $n \in \mathbb{N}$ with $\Pi_n(t)$ still being alive at time t. Note that $\Pi^x(t) \notin \mathcal{P}$ for each $t > 0$ as $\bigcup_{n\in\mathbb{N}} \Pi_n^x(t) \subsetneq \mathbb{N}$ due to the killing of blocks. Throughout this paper we shall repeatedly need an estimate regarding the number of fragments alive at a given random time $\tau : \Omega \to \mathbb{R}_0^+$. To this end, set $N_\tau^x := \operatorname{card}(\mathcal{N}_\tau^x)$ and observe that $N_\tau^x < \infty$ $\mathbb{P}$-almost surely. Indeed, since $\sum_{n\in\mathbb{N}} |\Pi_n(\tau)| \leq 1$ we infer that $|\Pi_n(\tau)| \geq e^{-(x+c\tau)}$ for at most $e^{x+c\tau}$-many $n \in \mathbb{N}$. Hence,

$$N_\tau^x \leq e^{x+c\tau}. \tag{3}$$

In addition, let ζ^x denote the random extinction time of Π^x, i.e. ζ^x is the supremum of all the killing times of individual blocks. The question whether ζ^x is finite or infinite was considered in [KK11]. Furthermore, define a function $\varphi : \mathbb{R} \to [0,1]$ by

$$\varphi(x) = \mathbb{P}(\zeta^x < \infty) \tag{4}$$

for all $x \in \mathbb{R}$. The function φ will be of utmost interest in the present paper. Let us point out that φ depends on the drift $c > 0$ of the killing line. However, in order to keep the notation as simple as possible, we omit this dependence in the notation as the constant c does not vary within results or proofs.

Set

$$\underline{p} := \inf\left\{p \in \mathbb{R} : \int_{\mathcal{S}} \left|1 - \sum_{n\in\mathbb{N}} s_n^{1+p}\right| \nu(\mathrm{d}\mathbf{s}) < \infty\right\} \in (-1, 0]$$

and for any $p > \underline{p}$ define

$$\Phi(p) := \int_{\mathcal{S}} \left(1 - \sum_{n\in\mathbb{N}} s_n^{1+p}\right) \nu(\mathrm{d}\mathbf{s})$$

as well as

$$c_p := \frac{\Phi(p)}{1+p}. \tag{5}$$

In view of [Ber03, Lemma 1] let $\bar{p}$ be the unique solution to the equation $(1+p)\Phi'(p) = \Phi(p)$ on $(\underline{p}, \infty)$, where Φ' denotes the derivative of Φ. In particular, this entails that $c_{\bar{p}} = \Phi'(\bar{p})$. According to [KK11] the killed fragmentation process survives with positive probability once the drift of the killing line is greater than $c_{\bar{p}}$ and becomes extinct almost surely otherwise.

Proposition 1 (Theorem 2 of [KK11]) *If $c > c_{\bar{p}}$, then $\varphi(x) \in (0,1)$ for all $x \in \mathbb{R}_0^+$. Otherwise, if $c \leq c_{\bar{p}}$, then $\varphi \equiv 1$.*

For any $t \in \mathbb{R}_0^+$ we denote by $B_n(t)$ the block of $\Pi(t)$ that contains the element $n \in \mathbb{N}$. According to Theorem 3 (ii) in [Ber01] it follows by means of the exchangeability of Π that under $\mathbb{P}$ the process

$$\xi_n := (-\ln(|B_n(t)|)\mathbb{1}_{\{|B_n(t)|>0\}})_{t\in\mathbb{R}_0^+},$$

cf. Figure 1(a), is a killed subordinator with Laplace exponent Φ, cemetery state $+\infty$ and killing rate

$$\int_{\mathcal{S}} \left(1 - \sum_{k\in\mathbb{N}} s_k\right) \nu(\mathrm{d}\mathbf{s}).$$

Hence, the process $X_n := (X_n(t))_{t\in\mathbb{R}_0^+}$, defined by

$$X_n(t) := ct - \xi_n(t)$$

for all $t \in \mathbb{R}_0^+$, is a spectrally negative Lévy process of bounded variation. Moreover, the jump times of X_n are precisely the times when the subordinator ξ_n jumps, which we denote by $(t_i)_{i\in\mathcal{I}_n}$. Note that the index set $\mathcal{I}_n$ is countably infinite. For $n \in \mathbb{N}$ and $x \in \mathbb{R}_0^+$ we shall make use of the shifted and killed process $X_n^x := (X_n^x(t))_{t\in\mathbb{R}_0^+}$, see Figure 1(b), given by

$$X_n^x(t) := (X_n(t) + x)\mathbb{1}_{\{\tau_{n,x}^- > t\}} = (x + ct + \ln(|B_n(t)|))\,\mathbb{1}_{\{\tau_{n,x}^- > t\}}$$

for every $t \in \mathbb{R}_0^+$, where

$$\tau_{n,x}^- := \inf\{t \in \mathbb{R}_0^+ : X_n(t) < -x\}.$$



(a) Illustration of Π, Π^x and ξ_n.

(b) Illustration of X_n^x.

Figure 1: Illustration (a) depicts Π, Π^x and ξ_n. The black dots indicate the blocks that belong to Π^x, whereas Π additionally contains the white dots. The dashed line constitutes the killing line with slope $c > 0$ starting at $x \in \mathbb{R}_0^+$. In addition, the bold piecewise constant graph illustrates the subordinator ξ_n. Notice that $\tau_{n,x}^-$ is the first time that ξ_n is above the killing line, i.e. it is the killing time of the block containing $n \in \mathbb{N}$. Illustration (b) shows the shifted and killed spectrally negative Lévy process X_n^x.

3 The one-sided FKPP equation for fragmentations

In this section we establish the set-up for our considerations by defining the FKPP travelling wave equation in the context of fragmentation processes. Moreover, the present section is devoted to presenting our main results. The main problem addressed in this paper is to find a regime of wave speeds for which we can prove the existence of a unique travelling wave solution of the one-sided FKPP equation for homogenous fragmentations as defined below. In order to tackle this problem we shall derive a theorem concerned with a product martingale that was introduced in [KK11]. Furthermore, we aim at studying some properties of travelling wave solutions.

3.1 Set-up

For any function $f : \mathbb{R} \to [0,1]$ set

$$\mathcal{C}_f := \left\{ x \in \mathbb{R}^+ : f'(x) \text{ exists} \right\}.$$

Since we do not know a priori that the functions f we are interested in are differentiable, we need to define the differential equations below for arguments in $\mathcal{C}_f$. We are particularly interested in the function φ, given by (4), and our main results in particular show that $\mathcal{C}_\varphi = \mathbb{R}^+$.

For certain functions $u : \mathbb{R}_0^+ \times \mathbb{R} \to [0,1]$, with $u|_{\mathbb{R}_0^+ \times (-\infty,0)} \equiv 1$ and $u(0,\cdot)|_{\mathbb{R}_0^+} = g$ for some continuous function $g : \mathbb{R}_0^+ \to [0,1]$, consider the initial value problem

$$\frac{\partial u}{\partial t}(t,x) = \int_{\mathcal{P}} \left(\prod_{n \in \mathbb{N}} u(t, x + \ln(|\pi_n|)) - u(t,x) \right) \mu(\mathrm{d}\pi) \tag{6}$$

for all $x \in \mathbb{R}_0^+$ and $t \in \mathcal{C}_{u(\cdot,x)}$. We call the above initial value problem *one-sided FKPP equation for fragmentation processes*. Here we are interested in the so-called FKPP travelling wave solutions of (6) with wave speed $c \in \mathbb{R}_0^+$, that is in solutions of (6) which are of the form $u(t,x) = f(x - ct)$ for all $t, x \in \mathbb{R}_0^+$.

Definition 2 A *one-sided FKPP travelling wave for fragmentation processes* is a monotone function $f : \mathbb{R} \to [0,1]$, with $f|_{(-\infty,0)} \equiv 1$, that satisfies the following *one-sided FKPP travelling wave equation*

$$cf'(x) + \int_{\mathcal{P}} \left(\prod_{n \in \mathbb{N}} f(x + \ln(|\pi_n|)) - f(x) \right) \mu(\mathrm{d}\pi) = 0 \tag{7}$$

for all $x \in \mathcal{C}_f$ with the boundary condition

$$\lim_{x \to \infty} f(x) = 0. \tag{8}$$

Let us now introduce an operator, whose definition is inspired by the integro-differential equation (7). To this end, let $\mathcal{D}_L$ be the set of all functions $f : \mathbb{R} \to [0,1]$, with $f|_{(-\infty,0)} \equiv 1$, for which the integral

$$\int_{\mathcal{P}} \left(\prod_{n \in \mathbb{N}} f(x + \ln(|\pi_n|) - f(x) \right) \mu(\mathrm{d}\pi)$$

exists, that is its positive part or negative part is finite. Then we define an integral operator L with domain $\mathcal{D}_L$ by

$$Lf(x) = \int_{\mathcal{P}} \left(\prod_{n \in \mathbb{N}} f(x + \ln(|\pi_n|) - f(x) \right) \mu(\mathrm{d}\pi)$$

for each $f \in \mathcal{D}_L$ and all $x \in \mathbb{R}_0^+$.

Let us remark that the upper Dini derivative (from above) of a function $f : \mathbb{R} \to [0,1]$ is defined by

$$f'_+(u) := \limsup_{h \downarrow 0} \frac{f(u+h) - f(u)}{h}$$

for all $u \in \mathbb{R}$. On this note, observe that this Dini derivative is well defined for any $u \in \mathbb{R}$, but may take the value ∞ or $-\infty$.

The following class of monotone functions plays a crucial role in the analysis of the one-sided FKPP travelling wave equation.

Definition 3 We denote by $\mathcal{T}$ the set of all nonincreasing functions $f : \mathbb{R} \to (0,1]$ with $f|_{(-\infty,0)} \equiv 1$ that satisfy (8) as well as

$$\sup_{u \in [s,t]} \underline{f}'_+(u) < \infty \tag{9}$$

for all $s, t \in \mathbb{R}_0^+$.

Note in particular that every $f \in \mathcal{T}$ is right-continuous on $\mathbb{R}^+$. Moreover, (9) implies that any $f \in \mathcal{T}$ admits an estimate of the form

$$f(x+h) - f(x) \leq |g(x+h) - g(x)| \tag{10}$$

for all $x \in \mathbb{R}^+$ and $h > 0$, where $g : \mathbb{R}^+ \to [0,1]$ is some continuous and right-differentiable function that has bounded right-derivatives on any compact set.

3.2 Main results

Here we present the main results of this paper. For this purpose, let us consider a product martingale which several of our proofs make use of. More precisely, for any function $f : \mathbb{R} \to [0,1]$ and $x \in \mathbb{R}_0^+$ let $Z^{x,f} := (Z_t^{x,f})_{t \in \mathbb{R}_0^+}$ be given by

$$Z_t^{x,f} = \prod_{n \in \mathcal{N}_t^x} f\left(X_n^x(t)\right)$$

for all $t \in \mathbb{R}_0^+$. This process was considered in Section 5 of [KK11] and in some proofs we shall resort to Theorem 10 in [KK11], which is concerned with the martingale property of $Z^{x,f}$. In this spirit, the first main result of the present paper reads as follows:

Theorem 1 *Let $c > c_{\bar{p}}$. In addition, let $f \in \mathcal{T}$ and assume that $Z^{x,f}$ is a martingale. Then f solves (7).*

The above theorem will be proven in Section 6. The following result, whose proof will be provided in Section 7, deals with some analytical properties of one-sided FKPP travelling waves.

Theorem 2 *Let $c > c_{\bar{p}}$. Then every one-sided FKPP travelling wave with wave speed c is an element of $\mathcal{T}$. Moreover, any such travelling wave is strictly monotonically increasing on $\mathbb{R}^+$ and continuously differentiable on $\mathbb{R}_0^+$, with a bounded derivative on each interval (x, ∞) with $x \in \mathbb{R}_0^+$.*

In particular, Theorem 2 shows that $-cf' = Lf$ on $\mathbb{R}^+$ for any one-sided FKPP travelling wave f with wave speed $c > c_{\bar{p}}$.

The main goal of this paper is to establish the existence of a unique travelling wave to (6) with wave speed c for $c > c_{\bar{p}}$ as well as the nonexistence of such a travelling wave with wave speed $c \leq c_{\bar{p}}$. More precisely, the following result states that the extinction probability of the killed fragmentation process solves equation (7) with boundary condition (8) for $c > c_{\bar{p}}$ and, moreover, is the only such function. Recall the function φ defined in (4).

Theorem 3 *If $c > c_{\bar{p}}$, then φ is the unique one-sided FKPP travelling wave with wave speed c. On the other hand, if $c \leq c_{\bar{p}}$, then there is no such travelling wave with wave speed c.*

We shall prove Theorem 3 in Section 8. In view of the forthcoming Remark 4 this theorem shows that one-sided FKPP travelling waves exist precisely for those positive wave speeds for which there do not exist two-sided travelling waves. Moreover, in view of [KK11, Theorem 4], Theorem 3 shows that travelling wave solutions exist exactly for those wave speeds that are larger than the asymptotic speed of the largest fragment in the killed fragmentation on the event of survival of this killed process. Let us further point out that if Π is conservative, i.e. loss of mass by sudden dislocations is excluded, then (7) can be written as

$$cf'(x) + \int_{\mathcal{S}} \left(\prod_{n \in \mathbb{N}} f(x + \ln(s_n)) - f(x) \right) \nu(\mathrm{d}\mathbf{s}) = 0,$$

which is the form of the two-sided FKPP equation considered in [BHK11] in the conservative setting, cf. (12).

4 Motivation – The classical FKPP equation

In order to present the framework in which Theorem 3 should be seen let us now briefly mention some known results that are related to our work. To this end we denote by $C^{1,2}(\mathbb{R}_0^+ \times A, [0,1])$, $A \subseteq \mathbb{R}$, the space of all functions $f : \mathbb{R}_0^+ \times A \to [0,1]$ such that $f(x,\cdot) \in C^2(A,[0,1])$ and $f(\cdot,y) \in C^1(\mathbb{R}_0^+,[0,1])$ for all $x \in \mathbb{R}_0^+$ and $y \in A$.

The classical FKPP equation in the form that is of most interest for us, cf. [McK75], is the following partial differential equation:

$$\frac{\partial u}{\partial t} = \frac{1}{2}\frac{\partial^2 u}{\partial x^2} + \beta(u^2 - u) \tag{11}$$

with $u \in C^{1,2}(\mathbb{R}_0^+ \times \mathbb{R}, [0,1])$. This equation, originally introduced by Fisher (see [Fis30] and [Fis37]) as well as by Kolmogorov, Petrovskii and Piscounov (cf. [KPP37]), has attracted much attention by analysts and probabilists alike. This kind of equation first arose in the context of a genetics model for the spread of an advantageous gene through a population and it is also satisfied by the maximal displacement of dyadic branching Brownian motion (see e.g. [McK75]). Several authors showed that this equation is closely related to dyadic branching Brownian motions, thus establishing a link of this analytical problem to probability theory. In this probabilistic interpretation the term "$\frac{1}{2}\frac{\partial^2 u}{\partial x^2}$" corresponds to the motion of the underlying Brownian motion, the "β" is the rate at which the particles split and the term "$u^2 - u$" results from the binary branching, that is two particles replace one particle at each branching time.

A solution u of equation (11) can be interpreted in different ways. The classical work concerning this partial differential equation, such as [Fis30], [Fis37] and [KPP37], describes the wave of advance of advantageous genes. More precisely, there are two types of individuals (or genes) in a population, and $u(t, x)$ measures the frequency or concentration of the advantageous type at the space-time point (x, t). In McKean's interpretation [McK75] the function $u(t, x)$ is related to a dyadic branching Brownian motion. Let $u(t, x)$ be the probability that at time t the largest particle of the branching Brownian motion has a value less than x. Then u satisfies equation (11). In [Fis30], [Fis37] and [KPP37] the equation describes the bulk of the population, in [McK75] it describes the most advanced particle.

The classical FKPP travelling waves are solutions of (11) of the form

$$u(t, x) = f(x - ct)$$

for some $f \in C^2(\mathbb{R}, [0, 1])$ and some constant $c \in \mathbb{R}$. This leads to the so-called FKPP travelling wave equation with wave speed $c \in \mathbb{R}$,

$$\begin{aligned} \frac{1}{2}f'' + cf' + \beta(f^2 - f) &= 0 \\ \lim_{x\to-\infty} f(x) &= 0 \\ \lim_{x\to\infty} f(x) &= 1, \end{aligned}$$

where $\beta > 0$. This travelling wave boundary value problem was studied by various authors, using both analytic as well as probabilistic techniques, and it is known that is has a unique (up to additive translation) solution $f \in C^2(\mathbb{R}, [0, 1])$ if $|c| \geq \sqrt{2\beta}$. In the opposite case that $0 \leq |c| < \sqrt{2\beta}$ there is no travelling wave solution. For probabilistic approaches we refer for instance to [McK75] (see also [McK76]), [Uch77], [Uch78], [Bra78], [Bra83], [Nev87], [CR88], [CR90] as well as [Har99] and [Kyp04].

Interesting with regard to our work is that the above boundary value problem was extended to continuous-time branching random walks (cf. [Kyp99]) and recently to conservative homogenous fragmentation processes (see [BHK11]). In the context of such fragmentation processes the corresponding partial integro-differential equation, referred to as FKPP equation, is given by

$$\frac{\partial u}{\partial t}(t, x) = \int_{\mathcal{S}} \left(\prod_{n\in\mathbb{N}} u(t, x + \ln(s_n)) - u(t, x) \right) \nu(\mathrm{d}\mathbf{s}) \tag{12}$$

for certain $u : \mathbb{R}_0^+ \times \mathbb{R} \to [0, 1]$. Note that (12) looks quite different compared to the classical FKPP equation (11). This difference results from the fact that fragmentation processes have no spatial motion except at jump times and from the more complicated jump structure of fragmentations in comparison with dyadic branching Brownian motions. However, despite the difference of the above equation compared to the classical FKPP equation, the name *FKPP equation* for (12) stems from the similarity in terms of the probabilistic interpretation these two equation have. Of particular interest are the FKPP travelling waves to (12) with wave speed $c \in \mathbb{R}$, i.e. solutions of (12) which are of the form $u(t, x) = f(x - ct)$ for all $t \in \mathbb{R}_0^+$ and $x \in \mathbb{R}$. These travelling wave solutions are functions $f \in C^1(\mathbb{R}, [0, 1])$ that satisfy the following FKPP travelling wave equation

$$cf'(x) + \int_{\mathcal{S}} \left(\prod_{n\in\mathbb{N}} f(x + \ln(s_n) - f(x) \right) \nu(\mathrm{d}\mathbf{s}) = 0$$

for all $x \in \mathbb{R}$ with boundary conditions

$$\lim_{x\to-\infty} f(x) = 0 \qquad \text{and} \qquad \lim_{x\to\infty} f(x) = 1.$$

Remark 4 For every $p \in (\underline{p}, \bar{p}]$ let $\mathcal{T}_2(p)$ denote the space of monotonically increasing functions $f \in C^1(\mathbb{R}, [0,1])$ satisfying the boundary conditions $\lim_{x\to-\infty} f(x) = 0$ as well as $\lim_{x\to\infty} f(x) = 1$ and such that $e^{(1+p)x}(1 - f(x))$ is monotonically increasing. In Theorem 1 of [BHK11] Berestycki et. al. showed that for $p \in (\underline{p}, \bar{p}]$ there exists a unique (up to additive translation) FKPP travelling wave solution in $\mathcal{T}_2(p)$ with wave speed c_p, cf. (5). According to Lemma 1 of [Ber03] the mapping $p \mapsto \Phi(p)/(1+p) = c_p$ is monotonically increasing on $(\underline{p}, \bar{p}]$ and thus it follows in view of Theorem 3 (ii) in [BHK11] that $c_{\bar{p}}$ is the maximal wave speed for two-sided travelling waves. $\Diamond$

In this paper we are interested in the one-sided counterpart of the abovementioned FKPP equation. In the classical setting the one-sided FKPP equation is the following partial differential equation

$$\frac{\partial u}{\partial t} = \frac{1}{2}\frac{\partial^2 u}{\partial x^2} + \beta(u^2 - u)$$

on $\mathbb{R}^+ \times \mathbb{R}^+$ with $u \in C^{1,2}(\mathbb{R}_0^+ \times \mathbb{R}_0^+, [0,1])$. Observe that this equation is the analogue of (11) for functions defined on $\mathbb{R}_0^+ \times \mathbb{R}_0^+$. The corresponding one-sided FKPP travelling wave equation with wave speed $c \in \mathbb{R}$ is given by the differential equation

$$\frac{1}{2}f'' + cf' + \beta(f^2 - f) = 0 \tag{13}$$

on $\mathbb{R}^+$ for $f \in C^2(\mathbb{R}_0^+, [0,1])$ satisfying the boundary conditions

$$\lim_{x\to 0} f(x) = 1 \qquad \text{as well as} \qquad \lim_{x\to\infty} f(x) = 0. \tag{14}$$

By considering killed branching Brownian motion with drift, killed upon hitting the origin, Harris et. al. proved in [HHK06] that solutions of the one-sided FKPP travelling wave boundary value problem (13) and (14) exist and are unique (up to additive translation) for all $c \in (-\sqrt{2\beta}, \infty)$ and there is no such travelling wave solution for $c \in (-\infty, -\sqrt{2\beta}]$. Notice that the one-sided travelling wave solutions for negative c are precisely those wave speeds for which there does not exist a two-sided travelling wave. With regard to the one-sided FKPP travelling wave equation in the classical setting we refer also to [Wat65], concerning existence of a solution, as well as [Pin95] for existence and uniqueness of a solution of (13) and (14) obtained by means of analytic techniques.

5 The finite activity case

In this section we prove existence of a one-sided FKPP travelling wave in the situation of a finite dislocation measure ν, sometimes referred to as the *finite activity case*. In this respect, note that even though a homogenous fragmentation process with finite ν may still have infinitely many jumps in any finite time interval, because infinitely many blocks may be present at any time and each block fragments with the same rate, in this setting every single block B_n has only finitely many jumps up to any $t \in \mathbb{R}_0^+$. In particular, this implies that the killed fragmentation with finite ν has finite activity in finite time intervals as at any time there are only finitely many blocks alive. Therefore, in this finite activity situation it is possible to consider the time $\tau_n : \Omega \to \mathbb{R}^+ \times \{\infty\}$ of the n-th jump of the killed process Π^x, a fact we shall make use of below.

An approach to solve the classical one-sided FKPP equation with boundary condition $u(0,x) = g(x)$ for some suitable function $g : \mathbb{R}_0^+ \to [0,1]$ is to show that the function $u : \mathbb{R}_0^+ \times \mathbb{R}_0^+ \to [0,1]$ given by

$$u(t,x) = \mathbb{E}\left(\prod_{n\in\mathbb{N}} g(x + Y_n(t))\right)$$

for all $t, x \in \mathbb{R}_0^+$ is a solution of the considered boundary value problem, where the $Y_n(t)$, $n \in \mathbb{N}$, are the positions of the particles at time t in a dyadic branching Brownian motion.

In this section we show that for fragmentations with a finite dislocation measure ν a similar approach as above works for the initial value problem (6). More precisely, we prove that for certain functions $g : \mathbb{R} \to [0,1]$ the function $u : \mathbb{R}_0^+ \times \mathbb{R} \to [0,1]$ defined by

$$u(t,x) = \mathbb{E}\left(\prod_{n \in \mathcal{N}_t^x} g(x + \ln(|B_n(t)|))\right) \qquad \text{and} \qquad u(t,\cdot)|_{(-\infty,0)} \equiv 1 \tag{15}$$

for all $t, x \in \mathbb{R}_0^+$ solves equation (6) with boundary condition $u(0,\cdot) = g$.

Proposition 5 *Assume that $\nu(\mathcal{S}) < \infty$ and let $c > 0$. Then every function $u : \mathbb{R}_0^+ \times \mathbb{R} \to [0,1]$ defined by (15), for some function $g : \mathbb{R} \to [0,1]$ with $g|_{\mathbb{R}_0^+} \in C^0(\mathbb{R}_0^+)$ and $g|_{(-\infty,0)} \equiv 1$, satisfies the boundary condition*

$$u(0,\cdot) = g \tag{16}$$

and solves equation (6) for any $x \in \mathbb{R}_0^+$ and Lebesgue-a.a. $t \in \mathbb{R}^+$. In particular, any such function u of the form $u(t,x) = f(x - ct)$ for some $f : \mathbb{R} \to [0,1]$, with $f|_{(-\infty,0)} \equiv 1$, and all $t \in \mathbb{R}_0^+$ and $x \in \mathbb{R}$ solves (7).

The following corollary of Proposition 5 provides a short proof that the extinction probability φ of the killed fragmentation process solves equation (7) in the special case of a finite dislocation measure.

Corollary 6 *Assume that $\nu(\mathcal{S}) < \infty$ and let $c > c_{\bar{p}}$. Then φ is an FKPP travelling wave with wave speed c.*

Proof Let us first show that φ solves (7). For this purpose, observe that the fragmentation property, in conjunction with the tower property of conditional expectations, yields that

$$\begin{aligned}\varphi(x - ct) &= \mathbb{E}\left(\prod_{n \in \mathcal{N}_t^x} \mathbb{P}\left(\zeta^{x-ct+ct+y} < \infty\right)\big|_{y=\ln(|B_n(t)|)}\right) \\ &= \mathbb{E}\left(\prod_{n \in \mathcal{N}_t^x} \varphi(x + \ln(|B_n(t)|))\right),\end{aligned}$$

and thus $u : \mathbb{R}_0^+ \times \mathbb{R} \to [0,1]$, given by $u(t,x) := \varphi(x - ct)$, satisfies (15) with $g = \varphi$. Hence, according to Proposition 5 the function φ solves (7). Since $c > c_{\bar{p}}$, it follows from Theorem 10 in [KK11] that φ also satisfies the boundary condition (8), which completes the proof. □

The major part of this paper, cf. Theorem 3, is concerned with the proof that the conclusion of Corollary 6 holds true also in the general case of an infinite dislocation measure.

Proof of Proposition 5 Let $g : \mathbb{R} \to [0,1]$ be some function that satisfies $g|_{\mathbb{R}_0^+} \in C^0(\mathbb{R}_0^+)$ and $g|_{(-\infty,0)} \equiv 1$. Further, consider the function $u : \mathbb{R}_0^+ \times \mathbb{R} \to [0,1]$ defined by (15) and fix some $t, x \in \mathbb{R}_0^+$. Note first that u satisfies the boundary condition (16), since $\mathcal{N}_0^x = \{1\}$ and $|B_1(0)| = 1$. In order to prove that u solves (6) Lebesgue-a.e., let $(t_i)_{i \in \mathcal{I}^x}$ be the jump times of Π^x and in view of the finiteness of the dislocation measure and $N_t^x \leq e^{x+ct}$, cf. (3), we assume without loss of generality that $\mathcal{I}^x = \mathbb{N}$ and that $0 < t_i < t_j$ for any $i, j \in \mathbb{N}$ with $i < j$. Since t_1 is exponentially distributed with parameter $\mu(\mathcal{P})$, we have

$$\lim_{h \downarrow 0} \frac{\mathbb{P}\left(t_1 \leq h\right)}{h} = \lim_{h \downarrow 0} \frac{1 - e^{-h\mu(\mathcal{P})}}{h} = \mu(\mathcal{P})$$

and deduce by resorting to the strong fragmentation property that

$$\lim_{h \downarrow 0} \frac{\mathbb{P}(t_2 \leq h)}{h} \leq \lim_{h \downarrow 0} \frac{\mathbb{P}(t_1 \leq h)^2}{h} = \lim_{h \downarrow 0} \frac{\left(1 - e^{-h\mu(\mathcal{P})}\right)^2}{h} = 0.$$

Consequently,

$$\lim_{h \downarrow 0} \frac{\mathbb{P}(t_1 \leq h, t_2 > h)}{h} = \lim_{h \downarrow 0} \frac{\mathbb{P}(t_1 \leq h)}{h} - \lim_{h \downarrow 0} \frac{\mathbb{P}(t_2 \leq h)}{h} = \mu(\mathcal{P}). \tag{17}$$

By means of the strong Markov property we have

$$\begin{aligned}
&\mathbb{E}\left(\prod_{n \in \mathcal{N}^x_{t+h}} g(x + \ln(|B_n(t+h)|))\mathbb{1}_{\left\{t_1^{(n)} - t \leq h, t_2^{(n)} - t > h\right\}}\right) \\
&= \mathbb{E}\left(\mathbb{E}\left(\prod_{n \in \mathcal{N}^x_{t+h}} g(x + \ln(|B_n(t+h)|))\mathbb{1}_{\left\{t_1^{(n)} - t \leq h, t_2^{(n)} - t > h\right\}} \middle| \mathscr{F}_t\right)\right) \\
&= \mathbb{E}\left(\prod_{k \in \mathcal{N}^x_t} \mathbb{E}\left(\prod_{n \in \mathbb{N}} g\left(x + \ln(y_k) + \ln\left(|\pi_n(t_1)|\right)\right)\right)\Bigg|_{y_k = |B_k(t)|}\right) \mathbb{P}(t_1 \leq h, t_2 > h) \\
&= \int_{\mathcal{P}} \prod_{n \in \mathbb{N}} \mathbb{E}\left(\prod_{k \in \mathcal{N}^x_t} g(x + \ln(|\pi_n|) + \ln(|B_k(t)|))\right) \frac{\mu(\mathrm{d}\pi)}{\mu(\mathcal{P})} \mathbb{P}(t_1 \leq h, t_2 > h) \\
&= \int_{\mathcal{P}} \prod_{n \in \mathbb{N}} u(t, x + \ln(|\pi_n|))\mu(\mathrm{d}\pi) \frac{\mathbb{P}(t_1 \leq h, t_2 > h)}{\mu(\mathcal{P})}.
\end{aligned}$$

where the $t_1^{(n)}$ and $t_2^{(n)}$ denote the first and second jump times respectively of the independent copy of Π^x that is initiated by the block $B_n(t)$. Therefore, (17) yields that

$$\begin{aligned}
&\lim_{h \downarrow 0} \mathbb{E}\left(\frac{1}{h} \prod_{n \in \mathcal{N}^x_{t+h}} g(x + \ln(|B_n(t+h)|))\mathbb{1}_{\left\{t_1^{(n)} - t \leq h, t_2^{(n)} - t > h\right\}}\right) \\
&= \int_{\mathcal{P}} \prod_{n \in \mathbb{N}} u(t, x + \ln(|\pi_n|))\mu(\mathrm{d}\pi) \lim_{h \downarrow 0} \frac{\mathbb{P}(t_1 \leq h, t_2 > h)}{\mu(\mathcal{P})} \\
&= \int_{\mathcal{P}} \prod_{n \in \mathbb{N}} u(t, x + \ln(|\pi_n|))\mu(\mathrm{d}\pi).
\end{aligned} \tag{18}$$

Moreover,

$$\mathbb{E}\left(\prod_{n \in \mathcal{N}^x_t} g(x + \ln(|B_n(t)|))\mathbb{1}_{\left\{t_1^{(n)} - t \leq h, t_2^{(n)} - t > h\right\}}\right)$$

$$
\begin{aligned}
&= \mathbb{E}\left(\prod_{n\in\mathcal{N}_t^x} g(x+\ln(|B_n(t)|))\mathbb{P}\left(t_1^{(n)}-t\le h, t_2^{(n)}-t>h \,\middle|\, \mathscr{F}_t\right)\right)\\
&= \int_{\mathcal{P}} u(t,x)\frac{\mu(\mathrm{d}\pi)}{\mu(\mathcal{P})}\mathbb{P}(t_1\le h, t_2>h)
\end{aligned}
$$

holds for all $h>0$ and hence

$$
\begin{aligned}
\lim_{h\downarrow 0}\mathbb{E}\left(\frac{1}{h}\prod_{n\in\mathcal{N}_t^x} g(x+\ln(|B_n(t)|))\mathbb{1}_{\left\{t_1^{(n)}-t\le h, t_2^{(n)}-t>h\right\}}\right) &= \int_{\mathcal{P}} u(t,x)\mu(\mathrm{d}\pi)\lim_{h\downarrow 0}\frac{\mathbb{P}(t_1\le h, t_2>h)}{\mu(\mathcal{P})}\\
&= \int_{\mathcal{P}} u(t,x)\mu(\mathrm{d}\pi). \qquad (19)
\end{aligned}
$$

Note that

$$
\prod_{n\in\mathcal{N}_{t+h}^x} g(x+\ln(|B_n(t+h)|))\mathbb{1}_{\left\{t_1^{(n)}-t>h\right\}} = \prod_{n\in\mathcal{N}_t^x} g(x+\ln(|B_n(t)|))\mathbb{1}_{\left\{t_1^{(n)}-t>h\right\}}
$$

for each $h>0$ as well as

$$
\lim_{h\downarrow 0}\frac{\mathbb{P}\left(t_2^{(n)}-t\le h\right)}{h} = 0.
$$

Since

$$
\begin{aligned}
&u(t+u,x)\\
&\mathbb{E}\left(\prod_{n\in\mathcal{N}_{t+u}^x} g(x+\ln(|B_n(t+u)|))\mathbb{1}_{\left\{t_1^{(n)}-t>h\right\}}\right)+\mathbb{E}\left(\prod_{n\in\mathcal{N}_{t+u}^x} g(x+\ln(|B_n(t+u)|))\mathbb{1}_{\left\{t_2^{(n)}-t\le h\right\}}\right)\\
&\quad+\mathbb{E}\left(\prod_{n\in\mathcal{N}_{t+u}^x} g(x+\ln(|B_n(t+u)|))\mathbb{1}_{\left\{t_1^{(n)}-t\le h, t_2^{(n)}-t>h\right\}}\right)
\end{aligned}
$$

holds for every $u\in\mathbb{R}_0^+$, it thus follows from (18) and (19) that

$$
\lim_{h\downarrow 0}\frac{u(t+h,x)-u(t,x)}{h} = \int_{\mathcal{P}}\left(\prod_{n\in\mathbb{N}} u(t,x+\ln(|\pi_n|))-u(t,x)\right)\mu(\mathrm{d}\pi),
$$

which completes the proof. □

6 Sufficiency criterion for existence of travelling waves

The goal of this section is to provide the proof of Theorem 1. In this regard, let us mention that Theorem 1 is a crucial tool for proving the other main results of the present paper. A first approach to try proving Theorem 1 might be to follow the lines of the proof of Theorem 1 in [BHK11]. But that proof relies on f being continuously differentiable and in our situation we cannot use any differentiability assumption. In fact, even if we knew that f is differentiable with a bounded derivative f', we would at least need that the set of discontinuities of f' is a Lebesgue null set. However, in general the set of such discontinuities may have positive Lebesgue measure, cf. Example 3.5 in [WH93].

We start by establishing two auxiliary results, which in spirit are analogues of respective results in [BHK11]. Afterwards we provide a lemma giving conditions under which only the block containing 1 is alive in the killed fragmentation process. Finally, having all these auxiliary results at hand, we finish this section with the proof of Theorem 1.

Observe first that a straightforward argument by induction yields that

$$\left|\prod_{n\in\mathbb{N}} a_n - \prod_{n\in\mathbb{N}} b_n\right| \le \sum_{n\in\mathbb{N}} |a_n - b_n| \tag{20}$$

for all sequences $(a_n)_{n\in\mathbb{N}}, (b_n)_{n\in\mathbb{N}} \in [0,1]^{\mathbb{N}}$.

Lemma 7 *Let $f \in \mathcal{T}$ and let $a, b \in \mathbb{R}_0^+$. Then*

$$\sup_{x\in[a,b]} \left|\prod_{n\in\mathbb{N}} f(x+\ln(|\pi_n|)) - f(x)\right| \in \mathscr{L}^1(\mu).$$

Proof By means of (20) we have

$$\begin{aligned}&\int_{\mathcal{P}} \sup_{x\in[a,b]} \left|\prod_{n\in\mathbb{N}} f(x+\ln(|\pi_n|)) - f(x)\right| \mu(\mathrm{d}\pi) \qquad (21)\\ &\le \int_{\mathcal{P}} \sup_{x\in[a,b]} \left|f(x+\ln(|\pi|_1^{\downarrow})) - f(x)\right| \mu(\mathrm{d}\pi) + \int_{\mathcal{P}} \sum_{n\in\mathbb{N}\setminus\{1\}} \sup_{x\in[a,b]} |f(x+\ln(|\pi_n|)) - 1|\, \mu(\mathrm{d}\pi).\end{aligned}$$

Since

$$\frac{\mathrm{d}}{\mathrm{d}x}[\ln(x) + 2(1-x)] = \frac{1}{x} - 2$$

and $\ln(1) + 2(1-1) = 0$, we deduce that

$$-\ln(x) \le 2(1-x)$$

holds for all $x \in [2^{-1}, 1]$. Therefore, for every $\epsilon \in (0, 2^{-1}]$ we have

$$-\ln(|\pi|_1^{\downarrow}) \le 2(1 - |\pi|_1^{\downarrow}) \tag{22}$$

for all $\pi \in \mathcal{P}$ with $1 - |\pi|_1^{\downarrow} \le \epsilon$. Moreover, by the Mean Value Theorem we have for any $x \in \mathbb{R}^+$ and $\pi \in \mathcal{P}$ with $|\pi|_1^{\downarrow} > e^{-x}$ the estimate

$$\left|f(x+\ln(|\pi|_1^{\downarrow})) - f(x)\right| \le \left|g(x+\ln(|\pi|_1^{\downarrow})) - g(x)\right| \le -\ln(|\pi|_1^{\downarrow}) \sup_{y\in(x+\ln(|\pi|_1^{\downarrow}),x)} |g'_+(y)|, \tag{23}$$

where the function g is given by (10). Notice that g'_+ denotes the right-derivative of g. Further, let $\epsilon \in (0, 2^{-1} \wedge (1-e^{-a}))$ and observe that $a + \ln(|\pi|_1^{\downarrow}) > 0$ for all $\pi \in \mathcal{P}$ with $1 - |\pi|_1^{\downarrow} < \epsilon$. Hence, resorting to (1), (22) and (23) we conclude that

$$\int_{\mathcal{P}} \sup_{x\in[a,b]} \left|f(x+\ln(|\pi|_1^{\downarrow})) - f(x)\right| \mu(\mathrm{d}\pi)$$

$$
\begin{aligned}
&\leq \int_{\{\pi\in\mathcal{P}:1-|\pi|_1^{\downarrow}\geq\epsilon\}} \sup_{x\in[a,b]} \left|f(x+\ln(|\pi|_1^{\downarrow}))-f(x)\right| \mu(\mathrm{d}\pi) \\
&\quad + \int_{\{\pi\in\mathcal{P}:1-|\pi|_1^{\downarrow}<\epsilon\}} \sup_{x\in[a,b]} \left|f(x+\ln(|\pi|_1^{\downarrow}))-f(x)\right| \mu(\mathrm{d}\pi) \\
&\leq \mu(\{\pi\in\mathcal{P}:1-|\pi|_1^{\downarrow}\geq\epsilon\}) + \sup_{y\in(a+\ln(|\pi|_1^{\downarrow}),b)} |g'_+(y)| \sup_{z\in[a,b]} \int_{\{\pi\in\mathcal{P}:1-|\pi|_1^{\downarrow}<\epsilon\}} -\ln(|\pi|_1^{\downarrow})\mu(\mathrm{d}\pi) \\
&\leq \mu(\{\pi\in\mathcal{P}:1-|\pi|_1^{\downarrow}\geq\epsilon\}) + 2 \sup_{y\in(a+\ln(|\pi|_1^{\downarrow}),b)} |g'_+(y)| \sup_{z\in[a,b]} \int_{\mathcal{P}} (1-|\pi|_1^{\downarrow})\mu(\mathrm{d}\pi) \\
&< \infty,
\end{aligned}
$$

which shows that the first term on the right-hand side of (21) is finite. In order to deal with the second term on the right-hand side of (21), note that the monotonicity of f together with $f|_{(-\infty,0)} \equiv 1$ and $f|_{[0,\infty)} \in [0,1]$ yields that

$$
\begin{aligned}
\int_{\mathcal{P}} \sum_{n\in\mathbb{N}\setminus\{1\}} \sup_{x\in[a,b]} |1-f(x+\ln(|\pi_n|))|\mu(\mathrm{d}\pi) &\leq \int_{\mathcal{P}} \sum_{n\in\mathbb{N}\setminus\{1\}} |1-f(b+\ln(|\pi_n|))|\mu(\mathrm{d}\pi) \\
&\leq \int_{\mathcal{P}} \sum_{n\in\mathbb{N}\setminus\{1\}} e^{(b+\ln(|\pi_n|))}\mu(\mathrm{d}\pi) \\
&= e^b \int_{\mathcal{P}} \sum_{n\in\mathbb{N}\setminus\{1\}} s_n\mu(\mathrm{d}\pi) \\
&< \infty
\end{aligned}
$$

for all $x > 0$. Observe that the finiteness follows from (1), since

$$
\int_{\mathcal{P}} \sum_{n\in\mathbb{N}\setminus\{1\}} s_n\mu(\mathrm{d}\pi) = \int_{\mathcal{P}} \left(\left(1-|\pi|_1^{\downarrow}\right) + \left(\sum_{n\in\mathbb{N}} s_n - 1\right)\right) \mu(\mathrm{d}\pi) \leq \int_{\mathcal{P}} (1-|\pi|_1^{\downarrow})\mu(\mathrm{d}\pi) < \infty.
$$

Consequently, we also have the finiteness of the second term on the right-hand side in (21). □

The next lemma shows that resorting to the previous lemma we obtain the continuity of Lf for every $f \in \mathcal{T}$.

Lemma 8 *Let $f \in \mathcal{T}$. Then the function Lf is continuous.*

Proof Fix some $x \in \mathbb{R}_0^+$ and let $(x_k)_{k\in\mathbb{N}}$ be a sequence in $\mathbb{R}^+$ with $x_k \to x$ as $k \to \infty$. Observe that for any $\epsilon > 0$ we have

$$
\begin{aligned}
&\int_{\mathcal{P}} \sup_{k\geq k_\epsilon} \left| \prod_{n\in\mathbb{N}} f(x+\ln(|\pi_n|)) - f(x) - \prod_{n\in\mathbb{N}} f(x_k+\ln(|\pi_n|)) + f(x_k)\right| \mu(\mathrm{d}\pi) \\
&\leq \int_{\mathcal{P}} \sup_{\delta\in[0,\epsilon]} \left| \prod_{n\in\mathbb{N}} f(x_{k_\epsilon}+\delta+\ln(|\pi_n|)) - f(x_{k_\epsilon}+\delta)\right| \mu(\mathrm{d}\pi) \qquad (24) \\
&\quad + 2\int_{\mathcal{P}} \sup_{\delta\in[0,\epsilon]} \left| \prod_{n\in\mathbb{N}} f(x+\delta+\ln(|\pi_n|)) - f(x+\delta)\right| \mu(\mathrm{d}\pi),
\end{aligned}
$$

where $k_\epsilon \in \mathbb{N}$ is chosen such that $|x - x_k| \leq |x - x_{k_\epsilon}| \leq \epsilon$ for all $k \geq k_\epsilon$. According to Lemma 7 both of the integrals on the right-hand side of (24) are finite. Hence, we can apply the DCT and deduce that

$$\begin{aligned}
&\lim_{k\to\infty} |Lf(x) - Lf(x_k)| \\
&= \int_{\mathcal{P}} \lim_{k\to\infty} \left| \prod_{n\in\mathbb{N}} f(x + \ln(|\pi_n|)) - f(x) - \prod_{n\in\mathbb{N}} f(x_k + \ln(|\pi_n|)) + f(x_k) \right| \mu(\mathrm{d}\pi) \\
&= \int_{\mathcal{P}} \left| \prod_{n\in\mathbb{N}} f(x + \ln(|\pi_n|)) - \prod_{n\in\mathbb{N}} \lim_{k\to\infty} f(x_k + \ln(|\pi_n|)) - f(x) + \lim_{k\to\infty} f(x_k) \right| \mu(\mathrm{d}\pi) \\
&= 0,
\end{aligned}$$

where the final equality follows from $f \in \mathcal{T}$ being continuous. Notice that we can interchange the limit and the product in the penultimate equality, since only finitely many factors of the product differ from 1. Hence, we have proven the continuity of Lf at x and since $x \in \mathbb{R}_0^+$ was chosen arbitrarily this completes the proof. $\square$

Recall that $(t_i)_{i\in\mathcal{I}_1}$ are the jump times of Π_1. For any $x \in \mathbb{R}_0^+$ consider the stopping time

$$\tau(x) := \inf\left\{ t \in (t_i)_{i\in\mathcal{I}_1} : |\pi(t)|_1^{\downarrow} \leq x \right\} \wedge 1.$$

Furthermore, observe that the assumption $\nu(\mathbf{s} \in \mathcal{S} : s_2 = 0) = 0$ ensures the existence of some $y \in (1/2, 1)$ such that $\nu(\mathbf{s} \in \mathcal{S} : s_1 \in (0, y]) > 0$.

Lemma 9 *Let* $x \in \mathbb{R}_0^+$ *and let* $y \in (1/2 \vee (1 - e^{-x}), 1)$ *be such that* $\nu(\mathbf{s} \in \mathcal{S} : s_1 \in (0, y]) > 0$. *Then*

$$\mathcal{N}_t^x \in \{\emptyset, \{1\}\}$$

$\mathbb{P}$-*a.s. on* $\{x + c\tau(y) < -\ln(1 - y)\}$ *for all* $t < \tau(y)$.

Lemma 9 says that up to time $\tau(y)$ the block in Π^x that contains 1 is the only block that may be alive on the event $\{x + c\tau(y) < -\ln(1 - y)\}$.

Proof As $y > 1/2$ we obtain

$$s_2 \leq 1 - s_1 < 1 - y < y \tag{25}$$

$\mathbb{P}$-a.s. for any $\mathbf{s} \in \mathcal{S}$ with $s_1 > y$. Moreover, since $|\pi(t_i)|_1^{\downarrow} > y$ for all $t_i < \tau(y)$, we have

$$-\ln(|\pi(t_i)|_2^{\downarrow}) \geq -\ln(1 - |\pi(t_i)|_1^{\downarrow}) \geq -\ln(1 - y) > x + c\tau(y) > x + ct_i$$

$\mathbb{P}$-a.s. on $\{x + c\tau(y) < -\ln(1 - y)\}$ for all $i \in \mathcal{I}_1^x$ with $t_i < \tau(y)$, where $(t_i)_{i\in\mathcal{I}_1^x}$ are the dislocation times of the block in Π^x containing 1. Notice that this implies that on $\{x + c\tau(y) < -\ln(1 - y)\}$ up to time $\tau(y)$ only the largest block can possibly survive as at each jump time before $\tau(y)$ the second-largest block in the resulting dislocation is killed instantaneously. However, bearing in mind the definition of $\tau(y)$ it follows from (25) that the block containing 1 is larger than the second-largest block resulting from any dislocation before time $\tau(y)$. Consequently, on the event $\{x + c\tau(y) < -\ln(1 - y)\}$ the block containing 1 is the only block that is possibly alive at some time $t \in [0, \tau(y))$. $\square$

We are now in a position to prove Theorem 1.

Proof of Theorem 1 Throughout the proof let $x \in \mathcal{C}_f$ and let $(a_n)_{n\in\mathbb{N}}$ be a $(0,1)$-valued sequence with $a_n \downarrow 0$ as $n \to \infty$.

The idea of the proof is to consider an appropriate decomposition of the limit of $\mathbb{E}(Z^{x,f}_{a_n} - Z^{x,f}_0)a_n^{-1}$ as $n \to \infty$, which by the martingale property of $Z^{x,f}$ equals 0. In this spirit, the proof is divided into three parts. The first and second part deal respectively with the jumps and drift that contribute to the difference $Z^{x,f}_{a_n} - Z^{x,f}_0$. In the third part we then combine the first two parts and resort to the martingale property of $Z^{x,f}$ in order to prove the assertion.

Part I Let us first deal with the jumps that contribute to the difference $Z^{x,f}_{a_n} - Z^{x,f}_0$. To this end, note first that the compensation formula for Poisson point processes yields that

$$\begin{aligned}
&\mathbb{E}\left(\sum_{i\in\mathcal{I}_1} \mathbb{1}_{\{t_i\in(0,a_n]\}}\left(\prod_{l\in\mathbb{N}} f(X^x_1(t_i-)+\ln(|\pi_l(t_i)|)) - f(X^x_1(t_i-))\right)\right)\\
&= \mathbb{E}\left(\int_{(0,a_n]}\int_{\mathcal{P}}\left(\prod_{l\in\mathbb{N}} f(X^x_1(t-)+\ln(|\pi_l|)) - f(X^x_1(t-))\right)\mu(\mathrm{d}\pi)\mathrm{d}t\right)\\
&= \mathbb{E}\left(\int_{(0,a_n]} Lf(X^x_1(t-))\mathrm{d}t\right).
\end{aligned}$$

Since

$$\min_{y\in[0,x+ca_n]} Lf(y) \le \mathbb{E}\left(\frac{1}{a_n}\int_{(0,a_n]} Lf(X^x_1(t-))\mathrm{d}t\right) \le \max_{y\in[0,x+ca_n]} Lf(y)$$

it thus follows in view of Lemma 8 that

$$\lim_{n\to\infty}\mathbb{E}\left(\frac{1}{a_n}\sum_{i\in\mathcal{I}_1} \mathbb{1}_{\{t_i\in(0,a_n]\}}\left(\prod_{l\in\mathbb{N}} f(X^x_1(t_i-)+\ln(|\pi_l(t_i)|)) - f(X^x_1(t_i-))\right)\right) = Lf(x). \tag{26}$$

In order to make the connection with the martingale $Z^{x,f}$ we shall consider the following the events $E_{\delta,t}$ and $E^{\complement}_{\delta,t}$ given by

$$E_{\delta,t} := \{\tau^-_{1,x}\wedge\tau(1-\delta) < t\} \cup \{\tau(1-\delta) \ge c^{-1}(-\ln(\delta)-x)\} \quad \text{and} \quad E^{\complement}_{\delta,t} := \Omega\setminus E_{\delta,t}$$

for all $\delta, t > 0$. Notice that the right-continuity of X_1 implies that $\tau^-_{1,x} > 0$ $\mathbb{P}$–almost surely. Moreover, according to Proposition 2 in Section 0.5 of [Ber96] the stopping time $\tau(1-\delta)$ is exponentially distributed and therefore $\mathbb{P}(\tau(1-\delta) > 0) = 1$. Consequently, since the map $\delta \mapsto c^{-1}(-\ln(\delta)-x)$ is decreasing and because $\tau(1-\delta)\downarrow 0$ as $\delta\downarrow 0$, we have

$$\lim_{\delta\downarrow 0}\lim_{n\to\infty}\mathbb{P}\left(E_{\delta,a_n}\right) = 0. \tag{27}$$

Observe that an application of the compensation formula results in

$$\begin{aligned}
&\frac{1}{a_n}\mathbb{E}\left(\sum_{i\in\mathcal{I}_1} \mathbb{1}_{\{t_i\in(0,a_n]\}}\mathbb{1}_{\{\tau^-_{1,x}\wedge\tau(1-\delta)\in(0,t_i)\}}\left(\prod_{l\in\mathbb{N}} f(X^x_1(t_i-)+\ln(|\pi_l(t_i)|)) - f(X^x_1(t_i-))\right)\right)\\
&= \frac{1}{a_n}\mathbb{E}\left(\int_{(0,a_n]}\mathbb{1}_{\{\tau^-_{1,x}\wedge\tau(1-\delta)\in(0,t)\}}\int_{\mathcal{P}}\left(\prod_{l\in\mathbb{N}} f(X^x_1(t_i-)+\ln(|\pi_l(t_i)|)) - f(X^x_1(t_i-))\right)\mu(\mathrm{d}\pi)\mathrm{d}t\right)
\end{aligned}$$

$$\leq \max_{y\in[0,x+ca_n]} Lf(y)\mathbb{P}\left(\tau_{1,x}^{-}\wedge\tau(1-\delta)<a_n\right) \tag{28}$$

Since $a_n \downarrow 0$ as $n\to\infty$, we infer from (27) and (28) that

$$\begin{aligned}&\mathbb{E}\left(\frac{1}{a_n}\sum_{i\in\mathcal{I}_1}\mathbb{1}_{\{t_i\in(0,a_n]\}}\mathbb{1}_{\{\tau_{1,x}^{-}\wedge\tau(1-\delta)\in(0,t_i)\}}\left(\prod_{l\in\mathbb{N}}f(X_1^x(t_i-)+\ln(|\pi_l(t_i)|))-f(X_1^x(t_i-))\right)\right)\\ &\quad\to 0 \end{aligned}\tag{29}$$

as $n\to\infty$ and then $\delta\downarrow 0$. Moreover, by means of the strong Markov property and the compensation formula we have

$$\begin{aligned}&\frac{1}{a_n}\mathbb{E}\left(\sum_{i\in\mathcal{I}_1}\mathbb{1}_{\{t_i\in(0,a_n]\}}\mathbb{1}_{\{\tau(1-\delta)\geq c^{-1}(-\ln(\delta)-x)\}}\left(\prod_{l\in\mathbb{N}}f(X_1^x(t_i-)+\ln(|\pi_l(t_i)|))-f(X_1^x(t_i-))\right)\right)\\ &\leq\frac{1}{a_n}\mathbb{E}\left(\sum_{i\in\mathcal{I}_1}\mathbb{1}_{\{t_i\in(0,a_n]\}}\left(\prod_{l\in\mathbb{N}}f(X_1^x(t_i-)+\ln(|\pi_l(t_i)|))-f(X_1^x(t_i-))\right)\mathbb{P}\left(\tau(1-\delta)\geq a_n+1|\,\mathscr{F}_{a_n}\right)\right)\\ &\leq\mathbb{E}\left(\frac{1}{a_n}\int_{(0,a_n]}\int_{\mathcal{P}}\left(\prod_{l\in\mathbb{N}}f(X_1^x(t-)+\ln(|\pi_l(t)|))-f(X_1^x(t-))\right)\mu(\mathrm{d}\pi)\mathrm{d}t\right)\mathbb{P}\left(\tau(1-\delta)\geq 1\right)\\ &\leq\max_{y\in[0,x+ca_n]}Lf(y)\mathbb{P}\left(\tau(1-\delta)\geq 1\right)\end{aligned}\tag{30}$$

for all $\delta\in(0,e^{-(x+2c)})$ which guarantees that $c^{-1}(-\ln(\delta)-x)\geq 2>a_n+1$. Since $\tau(1-\delta)\downarrow 0$ as $\delta\downarrow 0$, the estimates in (30) result in

$$\frac{1}{a_n}\mathbb{E}\left(\sum_{i\in\mathcal{I}_1}\mathbb{1}_{\{t_i\in(0,a_n]\}}\mathbb{1}_{\{\tau(1-\delta)\geq c^{-1}(-\ln(\delta)-x)\}}\left(\prod_{l\in\mathbb{N}}f(X_1^x(t_i-)+\ln(|\pi_l(t_i)|))-f(X_1^x(t_i-))\right)\right)\to 0$$

as $n\to\infty$ and then $\delta\downarrow 0$. Combining this with (26) we obtain

$$\lim_{\delta\downarrow 0}\lim_{n\to\infty}\mathbb{E}\left(\frac{1}{a_n}\sum_{i\in\mathcal{I}_1}\mathbb{1}_{\{t_i\in(0,a_n]\}}\left(\prod_{l\in\mathbb{N}}f(X_1^x(t_i-)+\ln(|\pi_l(t_i)|))-f(X_1^x(t_i-))\right)\mathbb{1}_{E_{\delta,t_i}}\right)=0$$

and in view of Lemma 9 we thus infer from (29) that

$$\begin{aligned}&\lim_{\delta\downarrow 0}\lim_{n\to\infty}\mathbb{E}\left(\frac{1}{a_n}\sum_{i\in\mathcal{I}_1}\mathbb{1}_{\{t_i\in(0,a_n]\}}\left(\prod_{l\in\mathbb{N}}f(X_1^x(t_i-)+\ln(|\pi_l(t_i)|))-f(X_1^x(t_i-))\right)\mathbb{1}_{E_{\delta,t_i}^{\complement}}\right)\\ &\quad=Lf(x).\end{aligned}\tag{31}$$

Furthermore, set

$$\Delta Z_t^{x,f}:=Z_t^{x,f}-Z_{t-}^{x,f}\tag{32}$$

for every $t>0$ and observe that f being bounded by 1, together with the strong Markov property and the compensation formula, implies by means of similar ideas as in (28) and (30) that

$$\frac{1}{a_n}\mathbb{E}\left(\sum_{i\in\mathcal{I}}\mathbb{1}_{\{t_i\in(0,a_n]\}}\Delta Z_{t_i}^{x,f}\mathbb{1}_{E_{\delta,t_i}}\right)$$

$$
\begin{aligned}
&\leq \frac{1}{a_n}\mathbb{E}\left(\sum_{i\in\mathcal{I}} \mathbb{1}_{\{t_i\in(0,a_n]\}}\mathbb{1}_{E_{\delta,t_i}}\left(\prod_{l\in\mathbb{N}} f(X^x_{j(t_i)}(t_i-)+\ln(|\pi_l(t_i)|)) - f(X^x_{j(t_i)}(t_i-))\right)\right)\\
&= \frac{1}{a_n}\mathbb{E}\left(\int_{(0,a_n]}\int_{\mathcal{P}} \mathbb{1}_{\{\tau^-_{1,x}\wedge\tau(1-\delta)\in(0,t)\}}\left(\prod_{l\in\mathbb{N}} f(X^x_{j(t)}(t-)+\ln(|\pi_l(t_i)|)) - f(X^x_{j(t)}(t-))\right)\mu(\mathrm{d}\pi)\mathrm{d}t\right)\\
&\quad+\frac{1}{a_n}\mathbb{E}\left(\int_{(0,a_n]}\int_{\mathcal{P}}\left(\prod_{l\in\mathbb{N}} f(X^x_{j(t)}(t-)+\ln(|\pi_l(t_i)|)) - f(X^x_{j(t)}(t-))\right)\mu(\mathrm{d}\pi)\mathrm{d}t\right)\mathbb{P}\left(\tau(1-\delta)\geq 1\right)\\
&\leq \max_{y\in[0,x+ca_n]} Lf(y)\left(\mathbb{P}\left(\tau^-_{1,x}\wedge\tau(1-\delta)<a_n\right)+\mathbb{P}\left(\tau(1-\delta)\geq 1\right)\right)
\end{aligned}
$$

for all $\delta\in(0,e^{-(x+2c)})$, where $j(t):=\inf\Pi^x_{k(t)}(t-)$, i.e. $B_{j(t)}(t-)=\Pi^x_{k(t)}(t-)$, for each $t>0$. Therefore, we obtain

$$
\lim_{\delta\downarrow 0}\lim_{n\to\infty}\mathbb{E}\left(\frac{1}{a_n}\sum_{i\in\mathcal{I}}\mathbb{1}_{\{t_i\in(0,a_n]\}}\Delta Z^{x,f}_{t_i}\mathbb{1}_{E_{\delta,t_i}}\right)=0. \tag{33}
$$

Consequently, since

$$
\begin{aligned}
&\mathbb{E}\left(\frac{1}{a_n}\sum_{i\in\mathcal{I}}\mathbb{1}_{\{t_i\in(0,a_n]\}}\Delta Z^{x,f}_{t_i}\mathbb{1}_{E^{\complement}_{\delta,t_i}}\right)\\
&\quad=\mathbb{E}\left(\frac{1}{a_n}\sum_{i\in\mathcal{I}_1}\mathbb{1}_{\{t_i\in(0,a_n]\}}\left(\prod_{l\in\mathbb{N}} f(X^x_1(t_i-)+\ln(|\pi_l(t_i)|))-f(X^x_1(t_i-))\right)\mathbb{1}_{E^{\complement}_{\delta,t_i}}\right),
\end{aligned}
$$

we deduce from (31) and (33) that

$$
\begin{aligned}
&\lim_{n\to\infty}\mathbb{E}\left(\frac{1}{a_n}\sum_{i\in\mathcal{I}}\mathbb{1}_{\{t_i\in(0,a_n]\}}\Delta Z^{x,f}_{t_i}\right)\\
&=\lim_{\delta\downarrow 0}\lim_{n\to\infty}\mathbb{E}\left(\frac{1}{a_n}\sum_{i\in\mathcal{I}}\mathbb{1}_{\{t_i\in(0,a_n]\}}\Delta Z^{x,f}_{t_i}\mathbb{1}_{E_{\delta,t_i}}\right)+\lim_{\delta\downarrow 0}\lim_{n\to\infty}\mathbb{E}\left(\frac{1}{a_n}\sum_{i\in\mathcal{I}}\mathbb{1}_{\{t_i\in(0,a_n]\}}\Delta Z^{x,f}_{t_i}\mathbb{1}_{E^{\complement}_{\delta,t_i}}\right)\\
&=Lf(x).
\end{aligned} \tag{34}
$$

Part II Let us now deal with the drift that contributes to the difference $Z^{x,f}_{a_n}-Z^{x,f}_0$. For this purpose, recall $\Delta Z^{x,f}_t$ from (32) and consider the process $(\hat{Z}^{x,f}_t)_{t\in\mathbb{R}^+_0}$, given by

$$
\hat{Z}^{x,f}_t := Z^{x,f}_t - \sum_{i\in\mathcal{I}^x:t_i\leq t}\Delta Z^{x,f}_{t_i}.
$$

Since on the event $E^{\complement}_{\delta,a_n}$ only the block containing 1 is alive, we only have to deal with X_1 on this event. Hence, since X_1 has drift c, we infer from (27) that

$$
\begin{aligned}
\lim_{\delta\downarrow 0}\lim_{n\to\infty}\mathbb{E}\left(\frac{\hat{Z}^{x,f}_{a_n}-\hat{Z}^{x,f}_0}{a_n}\mathbb{1}_{E^{\complement}_{\delta,a_n}}\right)&=\lim_{\delta\downarrow 0}\lim_{n\to\infty}\mathbb{E}\left(\frac{f(x+ca_n)-f(x)}{a_n}\mathbb{1}_{E^{\complement}_{\delta,a_n}}\right)\\
&=c\lim_{n\to\infty}\frac{f(x+ca_n)-f(x)}{ca_n}\lim_{\delta\downarrow 0}\lim_{n\to\infty}\mathbb{P}\left(E^{\complement}_{\delta,a_n}\right)
\end{aligned}
$$

$$= cf'(x). \tag{35}$$

Moreover, resorting to similar arguments as in (28) and (30) we deduce that

$$\begin{aligned}
&\frac{1}{a_n}\mathbb{E}\left(\sum_{i\in\mathcal{I}}\mathbb{1}_{\{t_i\in(0,a_n]\}}e^{x+ct_i}\mathbb{1}_{\{|\pi|_2^{\downarrow}\geq e^{-(x+ct_i)}\}}\mathbb{1}_{E_{\delta,t_i}}\right)\\
&\leq e^{x+ca_n}\mu\left(\pi\in\mathcal{P}:|\pi|_2^{\downarrow}\geq e^{-(x+ct_i)}\right)\mathbb{P}\left(\{\tau_{1,x}^{-}\wedge\tau(1-\delta)<a_n\}\cup\{\tau(1-\delta)\geq 1\}\right).
\end{aligned}$$

Consequently, since $N_t^x \leq e^{x+ca_n}$ for all $t\in[0,a_n]$ and because each process X_n, $n\in\mathbb{N}$, has drift c, we infer by means of (20) and the monotonicity of f that

$$\begin{aligned}
&\mathbb{E}\left(\left|\frac{\hat{Z}_{a_n}^{x,f}-\hat{Z}_0^{x,f}}{a_n}\right|\mathbb{1}_{E_{\delta,a_n}}\right)\\
&\leq \mathbb{E}\left(\frac{f(x)-f(x+ca_n)}{a_n}\mathbb{1}_{E_{\delta,a_n}}\right)+\mathbb{E}\left(\frac{1}{a_n}\sum_{i\in\mathcal{I}}\sum_{j\in\mathcal{N}_{t_i}^x}(1-f(x+ca_n)\mathbb{1}_{\{t_i\in(0,a_n]\}}\mathbb{1}_{E_{\delta,t_i}}\right)\\
&\leq c\,\frac{f(x)-f(x+ca_n)}{ca_n}\mathbb{P}\left(E_{\delta,a_n}\right)+\mathbb{E}\left(\sum_{i\in\mathcal{I}}\mathbb{1}_{\{t_i\in(0,a_n]\}}e^{x+ct_i}\mathbb{1}_{\{|\pi|_2^{\downarrow}\geq e^{-(x+ct_i)}\}}\mathbb{1}_{E_{\delta,t_i}}\right)\\
&\leq c\,\frac{f(x)-f(x+ca_n)}{ca_n}\mathbb{P}\left(E_{\delta,a_n}\right)\\
&\qquad+e^{x+ca_n}\mu\left(\pi\in\mathcal{P}:|\pi|_2^{\downarrow}\geq e^{-(x+ca_n)}\right)\mathbb{P}\left(\{\tau_{1,x}^{-}\wedge\tau(1-\delta)<a_n\}\cup\{\tau(1-\delta)\geq 1\}\right).
\end{aligned}$$

Since

$$\mu\left(\pi\in\mathcal{P}:|\pi|_2^{\downarrow}\geq e^{-(x+ct)}\right)\leq\mu\left(\pi\in\mathcal{P}:|\pi|_1^{\downarrow}\leq 1-e^{-(x+t)}\right)<\infty$$

for each $t\in\mathbb{R}_0^+$, we conclude in view of (27) that

$$\begin{aligned}
&\lim_{\delta\downarrow 0}\lim_{n\to\infty}\mathbb{E}\left(\left|\frac{\hat{Z}_{a_n}^{x,f}-\hat{Z}_0^{x,f}}{a_n}\right|\mathbb{1}_{E_{\delta,a_n}}\right)\\
&\leq e^{x}c\lim_{n\to\infty}\frac{f(x)-f(x+ca_n)}{ca_n}\lim_{\delta\downarrow 0}\lim_{n\to\infty}\mathbb{P}\left(E_{\delta,a_n}\right)\\
&\qquad+e^{x}\mu\left(\pi\in\mathcal{P}:|\pi|_2^{\downarrow}\geq e^{-x}\right)\lim_{\delta\downarrow 0}\lim_{n\to\infty}\mathbb{P}\left(\{\tau_{1,x}^{-}\wedge\tau(1-\delta)<a_n\}\cup\{\tau(1-\delta)\geq 1\}\right)\\
&=0.
\end{aligned}$$

By means of (35) this results in

$$\lim_{n\to\infty}\mathbb{E}\left(\frac{\hat{Z}_{a_n}^{x,f}-\hat{Z}_0^{x,f}}{a_n}\right)=cf'(x). \tag{36}$$

Part III Combining (34) with (36) yields that

$$0=\lim_{n\to\infty}\mathbb{E}\left(\frac{Z_{a_n}^{x,f}-Z_0^{x,f}}{a_n}\right)$$

$$= \lim_{n\to\infty} \mathbb{E}\left(\frac{1}{a_n}\sum_{i\in\mathcal{I}} \mathbb{1}_{\{t_i\in(0,a_n]\}}\Delta Z^{x,f}_{t_i}\right) + \lim_{n\to\infty}\mathbb{E}\left(\frac{\hat{Z}^{x,f}_{a_n} - \hat{Z}^{x,f}_0}{a_n}\right)$$
$$= Lf(x) + cf'(x)$$

holds for all $x \in \mathcal{C}_f$, where the first equality results from the martingale property of $Z^{x,f}$. Consequently f solves (7), which completes the proof. $\square$

7 Analytical properties of one-sided FKPP travelling waves

In this section we provide the proof of Theorem 2. Let us start with the following proposition that shows uniqueness of one-sided FKPP travelling waves with wave speed $c > c_{\bar{p}}$.

Below we shall resort to the following extension of the fundamental theorem of calculus to Dini derivatives, taken from [HT06].

Proposition 10 (Theorem 11 in [HT06]) *Let $a, b \in \mathbb{R}$ with $a < b$. If f is a continuous function that has a finite Dini derivative $f'_+(y)$ for all $y \in [a,b]$, then*

$$f(b) - f(a) = \int_{[a,b]} f'_+(y)\ dy, \tag{37}$$

provided that f'_+ is Lebesgue integrable over $[a,b]$.

This extended version of the fundamental theorem of calculus for Lebesgue integrals will be used in the proof of Proposition 11 that we are now going to present. Furthermore, we shall resort to Proposition 10 also in order to show in the proof of Theorem 2 that travelling waves are continuously differentiable. Let us emphasise that f having finite Dini derivatives is essential in Proposition 10. Indeed, for singular functions f, such as the Cantor function, the equality in (37) does not hold true as $f'_+ = 0$ Lebesgue-a.e., but f is not a constant function.

Proposition 11 *Any one-sided FKPP travelling wave f with wave speed $c > c_{\bar{p}}$ satisfies $f = \varphi$.*

Proof The method of proof is based upon ideas of the first part of the proof of Theorem 1 in [BHK11]. In order to use such ideas in our setting we shall make use of Proposition 10.

Let $f \in \mathcal{D}_L$ be a monotone function that solves (7) and (8). In addition, let $(x_n)_{n\in\mathbb{N}}$ be a sequence in $\mathbb{R}^+$ with $x_n \in \mathcal{C}_f$ for each $n \in \mathbb{N}$ and such that $x_n \downarrow 0$ as $n \to \infty$. Further, for any $n \in \mathbb{N}$ set $Z^{x,f,n}_t := \prod_{k\in\mathcal{N}^x_t} f(X^x_k(t) \vee x_n)$ and notice that the map

$$t \mapsto \hat{Z}^{x,f,n}_t := Z^{x,f,n}_t - \sum_{i\in\mathcal{I}^x : t_i \le t} \Delta Z^{x,f,n}_{t_i},$$

where $\Delta Z^{x,f,n}_t := Z^{x,f,n}_t - Z^{x,f,n}_{t-}$, is continuous. Moreover, for any $t \in \mathbb{R}^+_0$ the Dini derivative

$$Z^{x,f,n}_+(t) := \limsup_{h\downarrow 0} \frac{\hat{Z}^{x,f,n}_{t+h} - \hat{Z}^{x,f,n}_t}{h} = \sum_{i\in\mathcal{N}^x_t} cf'_+(X^x_i(t)\vee x_n) \prod_{j\in\mathcal{N}^x_t\setminus\{i\}} f(X^x_j(t)\vee x_n)$$
$$= \prod_{j\in\mathcal{N}^x_t} f(X^x_j(u)\vee x_n) \sum_{i\in\mathcal{N}^x_t} \frac{cf'_+(X^x_i(t)\vee x_n)}{f(X^x_i(t)\vee x_n)} \tag{38}$$

is Lebesgue measurable and, in view of (3), (9) and $f \in \mathcal{T}$, has finite expectation, since

$$Z_+^{x,f,n}(t) \leq \sum_{k \in \mathcal{N}_t^x} cf'_+(X_k^x(t) \vee x_n) \leq e^{x+ct} c \sup_{y \in [x_n, x+ct]} f'_+(y) < \infty.$$

Hence, according to Proposition 10 we obtain that

$$Z_t^{x,f,n} - Z_0^{x,f,n} = \hat{Z}_t^{x,f,n} - \hat{Z}_0^{x,f,n} + \sum_{i \in \mathcal{I}^x : t_i \leq t} \Delta Z_{t_i}^{x,f,n} = \int_{(0,t)} Z_+^{x,f,n}(s) \, \mathrm{d}s + \sum_{i \in \mathcal{I}^x : t_i \leq t} \Delta Z_{t_i}^{x,f,n} \tag{39}$$

holds true for all $t \in \mathbb{R}_0^+$. Further, define a function $\psi : \mathbb{R}_0^+ \times \mathcal{P} \times \mathbb{N} \to \mathbb{R}$ by

$$\psi(u, \pi, i) := \prod_{j \in \mathcal{N}_{u-}^x \setminus \{i\}} f(X_j^x(u-) \vee x_n) \prod_{k \in \mathbb{N}} f(X_i^x(u-) \vee x_n + \ln(|\pi_k|)) - \prod_{j \in \mathcal{N}_{u-}^x} f(X_j^x(u-) \vee x_n).$$

With η being the Poisson random measure on $\mathbb{R}_0^+ \otimes \mathcal{P} \otimes \mathbb{N}$ that determines Π, we deduce from (39), in conjunction with the compensation formula for Poisson point processes and Fubini's theorem that

$$\begin{aligned} &\mathbb{E}\left(Z_t^{x,f,n}\right) - \mathbb{E}\left(Z_0^{x,f,n}\right) \\ &= \mathbb{E}\left(\int_{(0,t)} Z_+^{x,f,n}(u) \, \mathrm{d}u\right) + \mathbb{E}\left(\int_{(0,t) \times \mathcal{P} \times \mathbb{N}} \psi(u, \pi, i) \eta(\mathrm{d}u, \mathrm{d}\pi, \mathrm{d}i)\right) \\ &= \mathbb{E}\left(\int_{(0,t)} Z_+^{x,f,n}(u) \, \mathrm{d}u\right) + \mathbb{E}\left(\int_{(0,t)} \int_{\mathbb{N}} \int_{\mathcal{P}} \psi(u, \pi, i) \mu(\mathrm{d}\pi) \sharp(\mathrm{d}i) \, \mathrm{d}u\right) \end{aligned} \tag{40}$$

for all $t \in \mathbb{R}_0^+$. Moreover, observe that

$$\begin{aligned} &\sum_{i \in \mathcal{N}_{u-}^x} \int_{\mathcal{P}} \psi(u, \pi, i) \mu(\mathrm{d}\pi) \\ &= \sum_{i \in \mathcal{N}_{u-}^x} \prod_{j \in \mathcal{N}_{u-}^x \setminus \{i\}} f(X_j^x(u-)) \int_{\mathcal{P}} \left(\prod_{k \in \mathbb{N}} f(X_i^x(u-) \vee x_n + \ln(|\pi_k|)) - f(X_i^x(u-) \vee x_n)\right) \mu(\mathrm{d}\pi) \\ &= \prod_{j \in \mathcal{N}_{u-}^x} f(X_j^x(u-) \vee x_n) \sum_{i \in \mathcal{N}_{u-}^x} \frac{1}{f(X_i^x(u-) \vee x_n)} Lf(X_i^x(u-) \vee x_n) \end{aligned} \tag{41}$$

$\mathbb{P}$-a.s. for every $u \in \mathbb{R}^+$. Observe that $f(X_i^x(\cdot))$ being monotone Lebesgue-a.e. for each $i \in \mathbb{N}$ implies that $f(X_i^x(\cdot))$ is differentiable Lebesgue-almost everywhere. By means of (38), (40) and (41) this results in

$$\begin{aligned} &\mathbb{E}\left(Z_t^{x,f,n}\right) - \mathbb{E}\left(Z_0^{x,f,n}\right) \\ &= \mathbb{E}\left(\int_{(0,t)} \prod_{j \in \mathcal{N}_u^x} f(X_j^x(u) \vee x_n) \sum_{i \in \mathcal{N}_u^x} \frac{1}{f(X_i^x(u) \vee x_n)} \left(cf'_+ + Lf\right)(X_i^x(u) \vee x_n) \mathrm{d}u\right) = 0 \end{aligned}$$

for all $n \in \mathbb{N}$ and $t \in \mathbb{R}_0^+$, where the final equality is a consequence of f satisfying (7). Letting $n \to \infty$ and resorting to the boundedness and continuity of f, the $\mathbb{P}$-a.s. finiteness of $\mathcal{N}_t^x$ as well as to the positivity of $X_j^x(t)$ for all $j \in \mathcal{N}_t^x$ this yields

$$\mathbb{E}\left(Z_t^{x,f}\right) = \mathbb{E}\left(Z_0^{x,f}\right) = f(x)$$

for every $t \in \mathbb{R}_0^+$. By means of the fragmentation property of Π we conclude that

$$\mathbb{E}\left(Z_{t+s}^{x,f} \middle| \mathscr{F}_t \right) = \prod_{n \in \mathcal{N}_t^x} \mathbb{E}\left(Z_s^{y,f} \right)\Big|_{y=X_n^x(t)} = \prod_{n \in \mathcal{N}_t^x} f(X_n^x(t)) = Z_t^{x,f}$$

$\mathbb{P}$-a.s. for all $s, t \in \mathbb{R}_0^+$. Consequently, $Z^{x,f}$ is a martingale and thus we deduce from Theorem 10 in [KK11] that $f = \varphi$. $\square$

Proposition 11 shows that in order to derive analytical properties of one-sided FKPP travelling waves we only need to consider the function φ. Bearing this in mind we proceed to prove Theorem 2. In order to obtain strict monotonicity of φ we shall use the following result.

Lemma 12 *Let $c > c_{\bar{p}}$. For any $0 \leq x < y < \infty$ there exists some $\alpha_{x,y} > 0$ such that*

$$\varphi(x) - \varphi(x+h) \geq \alpha_{x,y} \left(\varphi(y) - \varphi(y+h) \right)$$

for all $h > 0$.

Proof In the first part of this proof we show that for every deterministic time $t > 0$ the probability that X_1 reaches level $x > 0$ before time t is positive. In the second part we use this fact in order to obtain a lower bound of the probability that for some $n \in \mathbb{N}$ the process X_n^x hits a given level $y > x$ before some deterministic time $s > 0$. Subsequently we combine this lower bound with the estimate (3) of the number of blocks that are alive at a given time and with the positivity of the probability of extinction.

Part I According to Corollary 3.14 in [Kyp06] we have that $(\tau_{1,x})_{x \in \mathbb{R}_0^+}$ is a subordinator with either killing at an independent exponential *time* τ_e or with no killing in which case we set $\tau_e := \infty$. Moreover, by means of Proposition 1.7 in [Ber99] we thus infer that

$$\mathbb{P}\left(\tau_{1,x}^+ < t \right) = \mathbb{P}\left(\{ \tilde{\tau}_{1,x}^+ < t \} \cap \{ x < \tau_e \} \right) = \mathbb{P}\left(\tilde{\tau}_{1,x}^+ < t \right) \mathbb{P}\left(x < \tau_e \right) > 0 \tag{42}$$

holds for all $t > 0$, where $\tilde{\tau}_{1,x}^+$ is some non-killed subordinator, independent of τ_e, satisfying

$$\tilde{\tau}_{1,x}^+ \mathbb{1}_{\{x < \tau_e\}} = \tau_{1,x}^+ \mathbb{1}_{\{x < \tau_e\}}.$$

Let us now show that

$$\forall\, t > 0 : \, \mathbb{P}(\tau_{1,x}^+ < \tau_{1,0}^- \wedge t) > 0. \tag{43}$$

To this end, assume we had $\mathbb{P}(\tau_{1,x}^+ < \tau_{1,0}^- \wedge t_0) = 0$ for some $t_0 > 0$. Our goal is to show that this results in a contradiction. For this purpose, set $\tau_0^2 := 0$ and for every $n \in \mathbb{N}$ define

$$\tau_n^1 := \inf\{ t \geq \tau_{n-1}^2 : X_1(t) < 0 \},$$
$$\tau_n^2 := \inf\{ t \geq \tau_n^1 : X_1(t) = 0 \} \wedge t_0.$$

Recall that for X_1 the point 0 is irregular for $(-\infty, 0)$. Hence, there exists some $\delta > 0$ such that $\mathbb{P}(\tau_{1,0}^- \geq \delta) > 0$ and consequently we obtain by means of the strong Markov property that

$$\sum_{n \in \mathbb{N}} \mathbb{P}\left(\tau_n^1 - \tau_{n-1}^2 \geq \delta \,\middle|\, \mathscr{F}_{\tau_{n-1}^2} \right) = \sum_{n \in \mathbb{N}} \mathbb{P}\left(\tau_{1,0}^- \geq \delta \right) = \infty \tag{44}$$

$\mathbb{P}$-almost surely. Since $\{\tau_n^1 - \tau_{n-1}^2 \geq \delta\}$ is $\mathscr{F}_{\tau_n^2}$-measurable, we can apply an extended Borel-Cantelli lemma (see e.g. [Dur91, (3.2) Corollary in Chapter 4] or [Bre92, Corollary 5.29]) to deduce that

$$\left\{\{\tau_n^1 - \tau_{n-1}^2 \geq \delta\} \text{ happens infinitely often}\right\} = \left\{\sum_{n\in\mathbb{N}} \mathbb{P}\left(\tau_n^1 - \tau_{n-1}^2 \geq \delta \,\middle|\, \mathscr{F}_{\tau_{n-1}^2}\right) = \infty\right\}.$$

Thus (44) implies that $\tau_n^1 \to \infty$ $\mathbb{P}$-a.s. as $n \to \infty$, so in particular we have $\tau_n^1 > t_0$ for $n \in \mathbb{N}$ sufficiently large. An application of the strong Markov property therefore yields that

$$\mathbb{P}\left(\tau_{1,x}^+ < t_0\right) \leq \sum_{n\in\mathbb{N}} \mathbb{P}\left(\tau_{n,x}^3 < \tau_n^1 \wedge t_0 - \tau_{n-1}^2 \,\middle|\, \mathscr{F}_{\tau_{n-1}^2}\right) \leq \sum_{n\in\mathbb{N}} \mathbb{P}(\tau_{1,x}^+ < \tau_{1,0}^- \wedge t_0) = 0, \tag{45}$$

where

$$\tau_{n,x}^3 := \inf\{t \geq \tau_{n-1}^2 : X_1(t) > x\}$$

for all $n \in \mathbb{N}$. Since (45) contradicts (42), we conclude that (43) does indeed hold true.

Part II Let $0 \leq x < y < \infty$ and for any $t \in \mathbb{R}_0^+$ set $R_1^x(t) := \sup_{n\in\mathbb{N}} X_n^x(t)$. Then

$$\tau_y^+(x) := \inf\left\{t \in \mathbb{R}_0^+ : R_1^x(t) \geq y\right\}.$$

Note that $R_1^x(\tau_y^+(x)) = y$ if $\tau_y^+(x) < \infty$, since R_1^x does not jump upwards and thus creeps over the level y. Furthermore, let $s > 0$ and set $\gamma := e^{x+cs} - 1$ as well as

$$\alpha_{x,y} := \mathbb{P}\left(\tau_y^+(x) < \zeta^x \wedge s\right) \mathbb{P}(\zeta^y < \infty)^\gamma.$$

Observe that (43) and Proposition 1 imply that $\alpha_{x,y} > 0$, since

$$\mathbb{P}\left(\tau_y^+(x) < \zeta^x \wedge s\right) \geq \mathbb{P}\left(\tau_{1,y-x}^+ < \tau_{1,0}^- \wedge s\right).$$

By means of the strong fragmentation property we deduce that

$$\begin{aligned}
\varphi(x) - \varphi(x+h) &= \mathbb{P}(\zeta^x < \infty) - \mathbb{P}(\zeta^{x+h} < \infty) \\
&\overset{(*)}{\geq} \mathbb{P}\left(\tau_y^+(x) < \zeta^x \wedge s\right) \mathbb{P}(\zeta^y < \infty)^\gamma \left(\mathbb{P}(\zeta^y < \infty) - \mathbb{P}(\zeta^{y+h} < \infty)\right) \\
&= \alpha_{x,y} \left(\mathbb{P}(\zeta^y < \infty) - \mathbb{P}(\zeta^{y+h} < \infty)\right) \\
&= \alpha_{x,y} \left(\varphi(y) - \varphi(y+h)\right)
\end{aligned}$$

holds true for any $h > 0$, where the exponent γ in $(*)$ results from the estimate $N_s^x \leq e^{x+cs} = \gamma + 1$ $\mathbb{P}$-a.s., cf. (3). Notice that in $(*)$ we have used that the size of each block alive at time $\tau_y^+(x)$ is less than or equal to y as well as the monotonicity of the survival probability. ☐

Lemma 13 *Let $c > c_{\bar{p}}$. Then φ is strictly monotonically decreasing on $\mathbb{R}_0^+$.*

Proof Let $x \in \mathbb{R}_0^+$ and set

$$\gamma_x := \ln\left(|\pi(\zeta^x)|_1^{\downarrow} \left|\Pi_{k(\zeta^x)}^x(\zeta^x-)\right|\right).$$

According to Proposition 1 we have $\mathbb{P}(\zeta^x < \infty) > 0$ and hence

$$\mathbb{P}\left(\{\zeta^x < \infty\} \cap \bigcup_{n\in\mathbb{N}} \{x + c\zeta^x + \gamma_x \in (-n, 0)\}\right)$$

$$
\begin{aligned}
&= \mathbb{P}(\{\zeta^x < \infty\} \cap \{x + c\zeta^x + \gamma_x \in (-\infty, 0)\}) \\
&= \mathbb{P}(\zeta^x < \infty) \\
&> 0.
\end{aligned}
$$

Therefore, there exists some $z > 0$ such that

$$\mathbb{P}\left(\{\zeta^x < \infty\} \cap \{x + c\zeta^x + \ln(|\pi(\zeta^x)|_1^{\downarrow}) \in (-z, 0)\}\right) > 0$$

and thus the strong fragmentation property, in conjunction with Proposition 1, yields that

$$\mathbb{P}(\{\zeta^x < \infty\} \cap \{\zeta^{x+z} = \infty\}) \geq \mathbb{P}(\{\zeta^x < \infty\} \cap \{x + c\zeta^x + \gamma_x \in (-z, 0)\})\mathbb{P}(\zeta^0 = \infty) > 0.$$

Consequently, there exists some $z > 0$ such that

$$
\begin{aligned}
\mathbb{P}(\zeta^x < \infty) &= \mathbb{P}(\{\zeta^x < \infty\} \cap \{\zeta^{x+z} = \infty\}) + \mathbb{P}(\{\zeta^x < \infty\} \cap \{\zeta^{x+z} < \infty\}) \\
&> \mathbb{P}(\zeta^{x+z} < \infty), \qquad (46)
\end{aligned}
$$

where the final estimate follows from $\{\zeta^{x+z} < \infty\} \subseteq \{\zeta^x < \infty\}$.

Observe that (46) implies that for every $h > 0$ there exists some $y \geq x$ such that

$$\varphi(y) > \varphi(y + h). \tag{47}$$

Consequently, according to Lemma 12 there exists some $\alpha_{x,y} > 0$ such that

$$\varphi(x) - \varphi(x + h) \geq \alpha_{x,y}\,(\varphi(y) - \varphi(y + h)) > 0$$

holds for all $h > 0$ and $y \geq x$ satisfying (47). Since $x \in \mathbb{R}_0^+$ was chosen arbitrarily, this shows that φ is strictly monotonically decreasing. □

In the proof of Theorem 2 we shall make use of the theory of scale functions for spectrally negative Lévy processes. For this purpose, let W be the scale function of the spectrally negative Lévy process X_1. That is to say, W is the unique monotonically increasing function $W : \mathbb{R} \to \mathbb{R}_0^+$ which is equal to 0 on $(-\infty, 0)$, whose Laplace transform satisfies

$$\int_{(0,\infty)} e^{-\beta x} W(x)\, \mathrm{d}x = \frac{1}{\psi(\beta)},$$

where ψ denotes the Laplace exponent of X_1, and whose restriction to $\mathbb{R}_0^+$ is continuous.

Let us now tackle the proof of Theorem 2.

Proof of Theorem 2 Recall that we already proved the strict monotonicity of φ in Lemma 13. This proof is divided into two parts. In the first part we show that $\varphi \in \mathcal{T}$ and in the second part we prove that $\varphi|_{\mathbb{R}_0^+}$ is continuously differentiable. According to Proposition 11 this then completes the proof.

Part I Recall the definition of $\mathcal{T}$ in Definition 3. Observe first that φ is monotone and that Theorem 10 in [KK11] yields that φ satisfies (8). Hence, in order to prove $\varphi \in \mathcal{T}$ it remains to show that (9) holds. To this end, let $\mathfrak{n}$ denote the excursion measure of excursions $(\mathbf{e}_s)_{s \in \mathbb{R}_0^+}$ indexed by local time at the running maximum of the Lévy process X_1. Furthermore, for any such excursion

$\mathbf{e}$ let $\bar{\mathbf{e}}$ denote the height of this excursion. According to Lemma 8.2 in [Kyp06] the scale function W has a left-derivative on $\mathbb{R}^+$ given by

$$W'_-(x) = W(x)\mathfrak{n}(\bar{\mathbf{e}} \geq x)$$

for any $x \in \mathbb{R}^+$. Note that $\mathfrak{n}$ being σ-finite (cf. Theorem 6.15 in [Kyp06]) and monotone implies that

$$\sup_{y \geq x} \mathfrak{n}(\bar{\mathbf{e}} \geq y) = \mathfrak{n}(\bar{\mathbf{e}} \geq x) < \infty,$$

i.e. W has bounded left–derivatives on compact intervals. Moreover, we have

$$\mathbb{P}(\xi^x = \infty) \geq \mathbb{P}_x\left(\tau^+_{1,y} < \tau^-_{1,0}\right)\mathbb{P}(\xi^y = \infty)$$

for any $x, y \in \mathbb{R}^+$ with $x < y$, where under $\mathbb{P}_x$ the process X_1 is shifted to start in x. Therefore, we obtain

$$\varphi(x) - \varphi(x+h) \leq (1 - \varphi(x))\left(\frac{1}{\mathbb{P}_x\left(\tau^+_{1,h} < \tau^-_{1,0}\right)} - 1\right) \leq \frac{1}{\mathbb{P}_x\left(\tau^+_{1,h} < \tau^-_{1,0}\right)} - 1 \tag{48}$$

for any $h, x \in \mathbb{R}^+$. By means of (8.8) in Theorem 8.1 of [Kyp06] we have

$$\frac{1}{\mathbb{P}_x\left(\tau^+_{1,h} < \tau^-_{1,0}\right)} - 1 = \frac{W(x+h) - W(x)}{W(x)}. \tag{49}$$

Consequently, we deduce that $\varphi \in \mathcal{T}$.

Part II Note first that $\varphi|_{\mathbb{R}^+_0}$ is continuous. Indeed, by means of Lemma 8.6 in [Kyp06] we have that $W(0) > 0$, which in view of (48), (49) and the continuity of $W|_{\mathbb{R}^+_0}$ yields that $\varphi|_{\mathbb{R}^+_0}$ is continuous.

Assume that φ satisfies (7) on $\mathcal{C}_\varphi$. In view of the continuity of φ it follows from Lemma 8 that $L\varphi$ is continuous. Moreover, we deduce from (7) that $\varphi'_+ = -cL\varphi$ Lebesgue-a.e. on $\mathbb{R}^+$. Since the upper Dini derivative φ'_+ is bounded on any interval $[a, b] \subseteq \mathbb{R}^+$, it thus follows from Proposition 10 and Lebesgue's integrability criterion for Riemann integrals that

$$\varphi(b) - \varphi(a) = \int_a^b \varphi'_+(x)\ \mathrm{d}x = -\frac{1}{c}\int_a^b L\varphi(x)\ \mathrm{d}x = F(b) - F(a)$$

for all $a, b \in \mathbb{R}^+$, where F is an antiderivative of $-c^{-1}L\varphi$. Hence, we have $\varphi = F + \text{const.}$ on $\mathbb{R}^+$. Consequently, φ' exists on $\mathbb{R}^+$ and is continuous, that is $\varphi|_{\mathbb{R}^+_0} \in C^1(\mathbb{R}^+_0, [0, 1])$. □

8 Existence and uniqueness of one-sided travelling waves

This section is devoted to the proof of Theorem 3. Our method of proof makes use of the results that we established in the previous two sections.

Proof of Theorem 3 Part I of the proof shows the nonexistence of one-sided FKPP travelling waves for wave speeds $c \leq c_{\bar{p}}$ and Part II proves the existence and uniqueness of such travelling waves for wave speeds above the critical value $c_{\bar{p}}$.

Part I Fix some $c \leq c_{\bar{p}}$ as well as $x \in \mathbb{R}_0^+$ and let $f \in \mathcal{D}_L$ be a monotone function satisfying (7). Then the proof of Proposition 11 yields that $(Z_t^{x,f})_{t \in \mathbb{R}_0^+}$ is a uniformly integrable martingale and hence the $\mathbb{P}$–a.s. martingale limit $Z_\infty^{x,f} := \lim_{t\to\infty} Z_t^{x,f}$ satisfies

$$\mathbb{E}\left(Z_\infty^{x,f}\right) = \mathbb{E}\left(Z_0^{x,f}\right) = f(x). \tag{50}$$

Since $c \leq c_{\bar{p}}$, we have according to Proposition 1 that $\mathbb{P}(\zeta^x < \infty) = 1$, that is to say $\mathcal{N}_t^x \to \emptyset$ $\mathbb{P}$-a.s. as $t \to \infty$. Because the empty product equals 1, we thus infer that

$$Z_\infty^{x,f} = \lim_{t\to\infty} \prod_{n \in \mathcal{N}_t^x} f(X_n^x(t)) = 1$$

$\mathbb{P}$-almost surely. In view of (50) this implies that $f \equiv 1$. Consequently, there does not exist a monotone function $f \in \mathcal{D}_L$ that satisfies (7) and (8).

Part II Now let $c > c_{\bar{p}}$ and $x \in \mathbb{R}_0^+$. In the light of Proposition 11 it only remains to show that φ indeed is a travelling wave. Our goal is to apply Theorem 1. To this end, observe that the fragmentation property yields that

$$\varphi(x) = \mathbb{E}(\mathbb{P}(\zeta^x < \infty | \mathscr{F}_t)) = \mathbb{E}\left(\prod_{n \in \mathcal{N}_t^x} \varphi(X_n^x(t))\right) = \mathbb{E}\left(Z_t^{x,\varphi}\right).$$

By means of the fragmentation property we therefore deduce that

$$\mathbb{E}\left(Z_{t+s}^{x,\varphi}\middle|\, \mathscr{F}_t\right) = \prod_{n \in \mathcal{N}_t^x} \mathbb{E}\left(Z_s^{y,\varphi}\right)\big|_{y = X_n^x(t)} = \prod_{n \in \mathcal{N}_t^x} \varphi(X_n^x(t)) = Z_t^{x,\varphi}$$

holds $\mathbb{P}$-almost surely. Hence, $Z^{x,\varphi}$ is a $\mathbb{P}$-martingale. In the proof of Theorem 2 we have shown that $\varphi \in \mathcal{T}$, and consequently we infer from Theorem 1 that φ solves the integro-differential equation (7). This proves that φ is a one-sided FKPP travelling wave, since φ satisfies the boundary condition (8). The uniqueness was shown in Proposition 11 and thus the proof is complete. ☐

Acknowledgement

I would like to thank Andreas E. Kyprianou for the helpful discussions regarding the topic of this paper.

References

[BHK11] J. Berestycki, S. C. Harris, A. E. Kyprianou. Travelling waves and homogeneous fragmentation, *Ann. Appl. Probab.*, **21**, pp. 1749-1794, 2011

[Ber96] J. Bertoin. *Lévy Processes*, Cambridge University Press, 1996

[Ber99] J. Bertoin. *Subordinators: examples and applications.* Ecole d'été de Probabilités de Saint-Flour XXVII, Lecture Notes in Mathematics, Springer, **1717**, pp. 1-91, 1999

[Ber01] J. Bertoin. Homogeneous fragmentation processes, *Probab. Theory Related Fields*, **121**, pp. 301-318, 2001

[Ber03] J. Bertoin. The asymptotic behavior of fragmentation processes, *J. Europ. Math. Soc.*, **5**, pp. 395-416, 2003

[Ber06] J. Bertoin. *Random fragmentation and coagulation Processes*, Cambridge University Press, 2006

[Bra78] M. D. Bramson. Maximal displacement of branching Brownian motion, *Comm. Pure Appl. Math.*, **31** (5), pp. 531-581, 1978

[Bra83] M. D. Bramson. *Convergence of solutions of the Kolmogorov equation to travelling waves*, Mem. Amer. Math. Soc., 1983

[Bre92] L. Breiman. *Probability*, second edition, SIAM, 1992

[CR88] B. Chauvin, A. Rouault. KPP equation and supercritical branching Brownian motion in the subcritical speed area. Application to spatial

[CR90] B. Chauvin, A. Rouault. Supercritical branching Brownian motion and KPP equation in the critical speed area, *Math. Nachr.* **149**, pp. 41-59, 1990

[Dur91] R. Durrett. *Probability: theory and examples*, Duxbury Press, 1991

[Fis30] R. A. Fisher. *The genetical theory of natural selection*, Oxford University Press, 1930

[Fis37] R. A. Fisher. The wave of advance of advantageous genes, *Ann. Eugenics*, **7**, pp. 355-369, 1937

[HT06] J. W. Hagood, B. S. Thomson. Recovering a function from a Dini derivative, *Amer. Math. Monthly*, **113** (1), pp. 34-46, 2006

[HHK06] J.W. Harris, S. C. Harris, A. E. Kyprianou. Further probabilistic analysis of the Fisher-Kolmogorov-Petrovskii-Piscounov equation: one sided travelling waves, **42** (1), *Ann. Inst. H. Poincaré Probab. Statist.*, pp. 125-145, 2006

[Har99] S. C. Harris. Travelling-waves for the FKPP equation via probabilistic arguments, *Proc. Roy. Soc. Edinburgh Sect. A* **129** (3), pp. 503-517, 1999

[KK11] R. Knobloch, A. E. Kyprianou. Survival of homogenous fragmentation processes with killing, submitted, 2011, arXiv: 1104.5078

[KPP37] A. Kolmogorov, I. Petrovskii, N. Piscounov. Étude de l'équation de la diffusion avec croissance de la quantité de matière et son application à un problème biologique, *Bull. Moscow Univ., Math. Mech.*, **1** (1), pp. 1-25, 1937 [English translation by V. M. Volosov: A study of the diffusion equation with increase in the amount of substance, and its application to a biological problem, *Selected Works of A. N. Kolmogorov*, Vol. I, Kluwer, 1991]

[Kyp99] A. E. Kyprianou. A note on branching Lévy processes, *Stochastic Processes and their Applications*, **82** (1), pp. 1-14, 1999

[Kyp04] A. E. Kyprianou. Travelling wave solutions to the K-P-P equation: alternatives to Simon Harris' probabilistic analysis, *Ann. Inst. H. Poincaré Probab. Statist.*, **40**, pp. 53-72, 2004

[Kyp06] A. E. Kyprianou. *Introductory Lectures on Fluctuations of Lévy Processes with Applications*, Springer, 2006

[McK75] H. P. McKean. Application of Brownian motion to the equation of Kolmogorov-Petrovskii-Piskunov, *Comm. Pure Appl. Math.*, **28** (3), pp. 323-331, 1975

[McK76] H. P. McKean. A correction to "Application of Brownian motion to the equation of Kolmogorov-Petrovskii-Piskunov" (Comm. Pure Appl. Math.), *Comm. Pure Appl. Math.*, **29** (5), pp. 553-554, 1976

[Nev87] J. Neveu. Multiplicative martingales for spatial branching processes. *Seminar on Stochastic Processes* Progress in Probability and Statistics, Birkhäuser, **15**, pp. 223-242, 1987

[Pin95] R. G. Pinsky. K-P-P-type asymptotics for nonlinear diffusion in a large ball with infinite boundary data and on $\mathbb{R}^d$ with infinite initial data outside a large ball, *Comm. Part. Differ. Equat.*, **20** (7-8), pp. 1369-1393, 1995

[Uch77] K. Uchiyama. The behavior of solutions of the equation of Kolmogorov-Petrovsky-Piskunov, *Proc. Japan Acad. Ser. A Math. Sci.*, **53** (7), pp. 225-228, 1977

[Uch78] K. Uchiyama. The behavior of solutions of some non-linear diffusion equations for large time, *J. Math. Kyoto Univ.*, **18** (3), pp. 453-508, 1978

[Wat65] S. Watanabe. On the branching process for Brownian particles with an absorbing boundary, *J. Math. Kyoto Univ.*, **4**, pp. 385-398, 1965

[WH93] G. L. Wise, E. B. Hall. *Counterexamples in probability and real analysis*, Oxford University Press, 1993